



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cell Explorer™ Live Cell Tracking Kit

Blue Fluorescence

Ordering Information	Storage Conditions	Instrument Platform
Product Number: 22620 (2 plates)	Keep in freezer Protect from moisture and light	Fluorescence microscope Flow Cytometer

Introduction

Our Cell Explorer™ Live Cell labeling kits are a set of tools used to label cells for fluorescence microscopic investigations of cellular functions. The effective labeling of cells provides a powerful method for studying cellular events in a spatial and temporal context.

This particular kit is designed to uniformly label live cells in blue fluorescence for the studies that require the fluorescent tag molecules retained inside cells for a relatively longer time. The kit uses a non-fluorescent dye that carries a cell-retaining moiety. The dye becomes strongly fluorescent upon entering into live cells, and is trapped inside cells to give stable fluorescence signals. The dye is a hydrophobic compound that easily permeates intact live cells. The labeling process is robust, requiring minimal hands-on time. This Cell Explorer™ Live Cell labeling kit can be readily adapted for many different types of fluorescence platforms such as microplate assays, flow cytometry and fluorescence microscope. It is useful for a variety of studies, including cell adhesion, chemotaxis, multidrug resistance, cell viability, apoptosis and cytotoxicity. The kit provides all the essential components with an optimized cell-labeling protocol, and can be used for both proliferating and non-proliferating cells (either suspension or adherent cells).

Kit Components

Components	Amount
Component A: Track It™ Blue	2 vials
Component B: DMSO	1 vial (0.2 mL)
Component C: Assay Buffer	1 bottle (20 mL)

Assay Protocol

Brief Summary

Prepare samples → Remove the cell plate from incubator → Add 10 µL/well of 10X Track It™ Blue working solution → Stain the cells at RT for 15 minutes to 1 hour → Wash the cells → Examine the specimen under microscope at Ex/Em = 360/445 nm

Note: Thaw all the components to room temperature before opening.

1. Prepare Cells:

- 1.1 For adherent cells: Plate cells overnight in growth medium at 10,000 to 40,000 cells/well/90 µL for 96-well plates or 2,500 to 10,000 cells/well/20 µL for 384-well plates.
- 1.2 For non-adherent cells: Centrifuge the cells from the culture medium and then suspend the cell pellets in culture medium at 50,000-100,000 cells/well/90 µL for 96-well poly-D lysine plates or 10,000-25,000 cells/well/20 µL for 384-well poly-D lysine plates. Centrifuge the plates at 800 rpm for 2 minutes with brake off prior to the experiment.

Note: Each cell line should be evaluated on an individual basis to determine the optimal cell density.

2. Prepare Track It™ Blue stain solution:

- 2.1 Prepare 2 mM Track It™ Blue stock solution: Add 25 µL of DMSO (Component B) into one of the Track It™ Blue vials (Component A) to make 2 mM stock solution.

Note: The unused portion of the Track It™ stock solution should be stored at -20 °C. Avoid repeated freeze/thaw cycles.

2.2 Prepare 10X Track It™ Blue working solution: Dilute 2 mM of Track It™ Blue stock solution (from Step 2.1) into Assay Buffer (Component C) to make 5 to 50 μM Track It™ Blue working solution. The working solution should be prepared enough for all the wells at 10 μL /well with the appropriate concentration. For example, to get Track It™ Blue at the final concentration of 20 μM for one 96-well microplate, dilute 10 μL of the Track It™ Blue stock solution into 1 mL of Assay Buffer (Component C) to make 1 mL of 20 μM (10X) Track It™ Blue working solution.

Note 1: The final concentration of the Track It™ Blue should be empirically determined for different cell types and/or experimental conditions. It is recommended to test at the concentrations that are at least over a ten fold range.

Note 2: We found that 2 μM final in well concentration is sufficient for most of cell lines.

3. Stain the cells:

- 3.1 To the cell wells add 10X Track It™ Blue working solution (from Step 2.2), which should be equal to 1/10 of the volume of cell culture medium. For example, for a 96-well plate, add 10 μL /well of 10X Track It™ Blue working solution into the cells.
- 3.2 Incubate the cells in a 37 °C, 5% CO₂ incubator for 15 min to 1 hour.
- 3.3 Wash cells with Hanks and 20 mM HEPES buffer (HHBS) or an appropriate buffer.
- 3.4 Fill the cell wells with growth medium.
- 3.5 Analyze the cells using a fluorescence microscope or flow cytometer with DAPI filter sets (Ex/Em = 360/445 nm).




Figure 1. Image of U2OS cells stained with 2 μM Cell Explorer™ Live Cell Tracking Kit *Blue Fluorescence* in a Costar black wall/clear bottom 96-well plate.

References

1. Wolff M, Wiedenmann J, Nienhaus GU, Valler M, Heilker R. (2006) Novel fluorescent proteins for high-content screening. *Drug Discov Today*, 11, 1054.
2. Lee S, Howell BJ. (2006) High-content screening: emerging hardware and software technologies. *Methods Enzymol*, 414, 468.
3. Haasen D, Schnapp A, Valler MJ, Heilker R. (2006) G protein-coupled receptor internalization assays in the high-content screening format. *Methods Enzymol*, 414, 121.

Warning: This kit is only sold to end users. Neither resale nor transfer to a third party is allowed without written permission from AAT Bioquest. Chemical analysis of the kit components is strictly prohibited. Please call us at 408-733-1055 or e-mail us at info@aatbio.com if you have any questions.