



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-ssDNA/dsDNA [m3D8] Standard Size Ab00347-1.4

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

Isotype and Format: Mouse IgG1, Fc Silent™, Kappa

Clone Number: m3D8

Alternative Name(s) of Target: DNA; desoxyribonucleic acid; dsDNA; ssDNA

UniProt Accession Number of Target Protein:

Published Application(s): hydrololysis, SPR, ELISA

Published Species Reactivity:

Immunogen: 3`-biotin oligodeoxynucleotide ss(dN)40 with
N=CCATGAGTGATAACACTGCGGCCAACTTACTTCTGACAAC

Specificity: Binds unspecifically to double- and single-stranded DNA oligomers.

Application Notes: This catalytic antibody binds to DNA oligomers both single- and double-stranded, irrespective of the sequence, showing DNA-hydrolytic activity. DNA is the principle storage of genetic information and usually sequestered within the nucleus in Eukaryotes. Extracellular DNA is a key danger signal for the immune system and an antigen to disease-causing auto-antibodies (incl. DNA-hydrolysing antibodies) e.g. in Systemic Lupus Erythematosus. Antibody formats comprising different numbers of variable domains may be useful tools in investigating the significance of these catalytic antibodies in disease.

Antibody First Published in: Kim et al. 2006 Heavy and light chain variable single domains of an anti-DNA binding antibody hydrolyze both double- and single-stranded DNAs without sequence specificity. Journal of Biological Chemistry 2006; 281(22):15287-1595 [PMID:16551636](#)

Note on publication: Describes binding studies on mAB 3D8 on different types of oligos (ssDNA, dsDNA, dT:dA, dN:dN etc) using ELISA and SPR.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration:

1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.