



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Anti-S100A8 & S100A9 [5.5] Bulk Size Ab00815-1.1-BT

Developed in partnership with Ximbio ([www.ximbio.com](http://www.ximbio.com)).

**Isotype and Format:** Mouse IgG1, Kappa

**Clone Number:** 5.5

**Alternative Name(s) of Target:** MRP-8; MRP8; MRP-14; MRP14; CF antigen; CFAG; L1 molecule; p8,14; Calprotectin

**UniProt Accession Number of Target Protein:** P05109; P06702

**Published Application(s):** IP, WB, ELISA, FC, IHC

**Published Species Reactivity:** Human

**Immunogen:** BALB/c mice were immunized using cells from acute monocytic leukemia patients. The BALB/c spleen cell suspension was fused with Sp2/0-Ag14 cells, and mAb 5.5 was isolated from the hybridoma cells either by radioimmunoassay with <sup>125</sup>I-labelled F(ab')<sub>2</sub> fragments of sheep anti-mouse Ig as second layer, or by indirect immunofluorescence using rhodamine-conjugated goat anti-mouse Ig as the second layer.

**Specificity:** mAb 5.5 specifically recognises complexed MRP-8/MRP-14 via the hydrophobic residues in helix IV of MRP-8 (S100A8) exposed during heterodimer formation. mAb recognition of MRP-8 & MRP-14 is calcium dependent. 5.5 epitope is most likely to be formed by linear sequence Phe26-His27-Ala28 in calcium-binding loop I of MRP-8, and by Phe68, Leu72, Ile76 and Lys77 that lie along one face of helix IV.

**Application Notes:** mAb 5.5 has been used in immunoprecipitation and western blot experiments for the characterisation of p8,14 identified in extracts of monocytes and neutrophils. 5.5 was also used in flow cytometric analysis to show that p8,14 is not a cell surface protein, and in immunohistochemical staining to show that p8,14 is present extracellularly (Edgeworth 1991).

**Antibody First Published in:** Hogg et al. Monoclonal antibody 5.5 reacts with p8, 14, a myeloid molecule associated with some vascular endothelium 1989. Eur J Immunol. 19(6):1053-61 [PMID:2666142](#)

**Note on publication:** Describes the activation of Myeloid cells by inflammatory stimuli, resulting in the secretion of p8,14 onto the endothelium. 5.5 was used for the detection of p8,14 to suggest that this molecule is released by myeloid cells adhering onto the endothelium, rather than the endothelium itself in response to inflammatory stimuli.

## Product Form

**Size:** 1 mg Purified antibody in bulk size.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommended this antibody be handled under sterile conditions. For longer storage, aliquot and store at -20°C.

**Concentration:** 1 mg/ml.

Important note - This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.