

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com




FLT1 & PTPN11 Protein Protein Interaction Antibody Pair

Catalog # : DI0015

規格 : [1 Set]

List All




Reactivity:

ity: Human

Quality Control Protein protein interaction immunofluorescence result.

Testing:




	20°C storage immediately after use.
MSDS:	Download
Publication Ref	erence
novel prognos Liu CH, Chen Cheng HC, Cl Proteomics. 2	f protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma. TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, nen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 013 Feb 8. [Epub ahead of print]
Applications	Ligation Apony (Call)
<i>In situ</i> Proximity	Ligation Assay (Cell)
FLT1 PTPN11	
Gene Informatio	on
Entrez GenelD:	2321
Gene Name:	FLT1
Gene Alias:	FLT,VEGFR1
Gene Description:	fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor)
Omim ID:	<u>165070</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (lg)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia
Other Designations:	fms-related tyrosine kinase 1,soluble VEGF receptor 1-14,soluble VEGFR1 variant 2,soluble VEGFR1 variant 21,vascular endothelial growth factor/vascular permeability factor receptor
Gene Informatio	on
Entrez GenelD:	5781
Gene Name:	PTPN11
Gene Alias:	BPTP3,CFC,MGC14433,NS1,PTP-1D,PTP2C,SH-PTP2,SH-PTP3,SHP2
Gene Description:	protein tyrosine phosphatase, non-receptor type 11
Omim ID:	<u>151100, 163950, 176876, 607785</u>
Gene Ontology	<u>Hyperlink</u>


Gene Summary: The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains two tandem Src homology-2 domains, which function as phospho-tyrosine binding domains and mediate the interaction of this PTP with its substrates. This PTP is widely expressed in most tissues and plays a regulatory role in various cell signaling events that are important for a diversity of cell functions, such as mitogenic activation, metabolic control, transcription regulation, and cell migration. Mutations in this gene are a cause of Noonan syndrome as well as acute myeloid leukemia. [provided by RefSeq

Otherprotein tyrosine phosphatase-2,protein-tyrosine phosphatase 2CDesignations:

Interactome 1



Interactome 2



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.