

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





TGFB1 & APP Protein Protein Interaction Antibody Pair

Catalog #: DI0054 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the TGFB1 protein, and the other against the APP protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




1. Incubate with target primary antibodies




2. Add PLA probes PLUS and MINUS



3. Hybridize connector oligos



4. Ligation to form a complete DNA circle



5. Rolling circle amplification




6. Add fluorescent probes to reveal interaction

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.

Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between TGFB1 and APP. HeLa cells were stained with anti-TGFB1 rabbit purified polyclonal antibody 1:1200 and anti-APP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied **Product:**

Antibody pair set content:

- 1. TGFB1 rabbit purified polyclonal antibody (20 ug)
- 2. APP mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -



Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

APP TGFB1

Gene Information

Entrez GeneID: 7040

Gene Name: TGFB1

Gene Alias: CED, DPD1, TGFB, TGFbeta

Gene transforming growth factor, beta 1

Description:

Omim ID: <u>131300</u>, <u>190180</u>, <u>219700</u>

Gene Ontology: Hyperlink

Gene Summary: TGFB is a multifunctional peptide that controls proliferation,

differentiation, and other functions in many cell types. TGFB acts synergistically with TGFA (MIM 190170) in inducing transformation. It also acts as a negative autocrine growth factor. Dysregulation of TGFB activation and signaling may result in apoptosis. Many cells synthesize TGFB and almost all of them have specific receptors for this peptide. TGFB1, TGFB2 (MIM 190220), and TGFB3 (MIM 190230) all function through the same receptor signaling systems.[supplied by OMIM

Other TGF-beta 1 protein, diaphyseal dysplasia 1, progressive, transforming

Designations: growth factor-beta 1

Gene Information

Entrez GenelD: 351

Gene Name: APP

Gene Alias: AAA,ABETA,ABPP,AD1,APPI,CTFgamma,CVAP,PN2

Gene amyloid beta (A4) precursor protein

Description:

Omim ID: <u>104760</u>, <u>605714</u>

Gene Ontology: Hyperlink

Gene Summary: This gene encodes a cell surface receptor and transmembrane


precursor protein that is cleaved by secretases to form a number of peptides. Some of these peptides are secreted and can bind to the acetyltransferase complex APBB1/TIP60 to promote transcriptional activation, while others form the protein basis of the amyloid plaques found in the brains of patients with Alzheimer disease. Mutations in this

gene have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid angiopathy). Multiple transcript variants encoding several different isoforms have been found for this gene. [provided by RefSeq


Other Designations:

A4 amyloid protein,amyloid beta A4 protein,amyloid-beta protein,beta-amyloid peptide,cerebral vascular amyloid peptide,peptidase nexin-II,protease nexin-II

Interactome 1



Interactome 2



Page 3 of 4 2016/5/19

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 4 of 4 2016/5/19