

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





## **BUB1 & EIF4EBP1 Protein Protein Interaction Antibody Pair**

Catalog #: DI0156 規格:[1 Set]

#### List All

### **Specification**

# Product Description:

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the BUB1 protein, and the other against the EIF4EBP1 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

## **Application Image**


In situ Proximity Ligation Assay (Cell)



 Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



3. Hybridize connector oligos



 Ligation to form a complete DNA circle



5. Rolling circle amplification




Add fluorescent probes
to reveal interaction

Reactivity:

Human

## Quality Contro Testing:

Quality Control Protein protein interaction immunofluorescence result.



Representative image of Proximity Ligation Assay of protein-protein interactions between BUB1 and EIF4EBP1. HeLa cells were stained with anti-BUB1 rabbit purified polyclonal antibody 1:1200 and anti-EIF4EBP1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

# Supplied Product:

Antibody pair set content:

- 1. BUB1 rabbit purified polyclonal antibody (20 ug)
- 2. EIF4EBP1 mouse monoclonal antibody (40 ug)

\*Reagents are sufficient for at least 30-50 assays using recommended protocols.

### Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

20°C storage immediately after use.

MSDS:

**Download** 

#### **Publication Reference**

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Chen JHO, Chen MH, Chen CF, Tian YO, Kan CY, Hairan M, Lhann CY, Mal Coll

Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

### **Applications**

In situ Proximity Ligation Assay (Cell)

BUB1 EIF4EBP1

**Gene Information** 

Entrez GenelD: 699

Gene Name: BUB1

Gene Alias: BUB1A,BUB1L,hBUB1

Gene budding uninhibited by benzimidazoles 1 homolog (yeast)

**Description:** 

Omim ID: <u>602452</u>

Gene Ontology: Hyperlink

**Gene Summary:** This gene encodes a kinase involved in spindle checkpoint function.

The kinase functions in part by phosphorylating a member of the miotic checkpoint complex and activating the spindle checkpoint. Mutations in this gene have been associated with aneuploidy and several forms of

cancer. [provided by RefSeq

Other BUB1 budding uninhibited by benzimidazoles 1 homolog, budding

**Designations:** uninhibited by benzimidazoles 1,mitotic spindle checkpoint

kinase, putative serine/threonine-protein kinase

**Gene Information** 

Entrez GeneID: 1978

Gene Name: EIF4EBP1

Gene Alias: 4E-BP1,4EBP1,BP-1,MGC4316,PHAS-I

Gene eukaryotic translation initiation factor 4E binding protein 1

Description:

Omim ID: <u>602223</u>

Gene Ontology: Hyperlink

Gene Summary: This gene encodes one member of a family of translation repressor

proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in


response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA

translation. [provided by RefSeq


Other elF4E-binding protein 1,phosphorylated heat- and acid-stable protein

**Designations:** regulated by insulin 1

### Interactome 1



### Interactome 2



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/19