

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



## **TP53 & HSPA1B Protein Protein Interaction Antibody Pair**

Catalog # : DI0230

規格 : [1 Set]

List All


Specification				Application Image
Product Description:	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the TP53 protein, and the other against the HSPA1B protein for use in <u>in</u> <u>situ Proximity Ligation Assay</u> . See Publication Reference below.			<i>In situ</i> Proximity Ligation Assay (Cell)
	1. Incubate with target primary antibodies	2. Add PLA probes PLUS and MINUS	3. Hybridize connector oligos	
	4. Ligation to form a complete DNA circle	5. Rolling circle amplification	6. Add fluorescent probes to reveal interaction	

Reactivity: Human

.....

**Quality Control** Protein protein interaction immunofluorescence result.

Testing:




Representative image of Proximity Ligation Assay of protein-protein interactions between TP53 and HSPA1B. HeLa cells were stained with anti-TP53 rabbit purified polyclonal antibody 1:1200 and anti-HSPA1B mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (<u>BlobFinder</u>) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:	Antibody pair set content: 1. TP53 rabbit purified polyclonal antibody (20 ug) 2. HSPA1B mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

	20°C storage immediately after use.
MSDS:	m Download
Publication Ref	erence
novel prognos Liu CH, Chen Cheng HC, Cł Proteomics. 2	f protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma. TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, hen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 013 Feb 8. [Epub ahead of print]
Applications In situ Proximity	/ Ligation Assay (Cell)
HSPA1B TP53	
Gene Informatio	
Entrez GenelD:	
Gene Name:	
Gene Alias:	FLJ92943,LFS1,TRP53,p53
Gene Description:	tumor protein p53
Omim ID:	<u>114480, 114500, 114550, 151623, 161550, 191170, 202300, 260350</u>
Gene Ontology:	: <u>Hyperlink</u>
Gene Summary:	This gene encodes tumor protein p53, which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of this gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome. Multiple p53 variants due to alternative promoters and multiple alternative splicing have been found. These variants encode distinct isoforms, which can regulate p53 transcriptional activity. [provided by RefSeq
Other Designations:	p53 antigen,p53 transformation suppressor,p53 tumor suppressor,phosphoprotein p53,transformation-related protein 53
Gene Informatio	วท
Entrez GenelD:	3304
Gene Name:	HSPA1B
Gene Alias:	FLJ54328,HSP70-1B,HSP70-2,HSPA1A

Omim ID:	603012	
Gene Ontology: <u>Hyperlink</u>		
·	This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjuction with other heat shock proteins, this protein stabilizes existing proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility complex class III region, in a cluster with two closely related genes which encode similar proteins. [provided by RefSeq	
	OTTHUMP00000029288,heat shock 70kD protein 1B,heat shock 70kDa protein 1A	



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.