

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





EGFR & EGF Protein Protein Interaction Antibody Pair

Catalog #: DI0240 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the EGFR protein, and the other against the EGF protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




 Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



Hybridize connector oligos



 Ligation to form a complete DNA circle



Rolling circle amplification




Add fluorescent probes
to reveal interaction

Reactivity:

Human

Quality Contro Testing:

Quality Control Protein protein interaction immunofluorescence result.



Representative image of Proximity Ligation Assay of protein-protein interactions between EGFR and EGF. HeLa cells were stained with anti-EGFR rabbit purified polyclonal antibody 1:1200 and anti-EGF mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:

Antibody pair set content:

- 1. EGFR rabbit purified polyclonal antibody (20 ug)
- 2. EGF mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

20°C storage immediately after use.

MSDS:

<u>m</u>Download

Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

EGF EGFR

Gene Information

Entrez GenelD: 1956

Gene Name: EGFR

Gene Alias: ERBB,ERBB1,HER1,PIG61,mENA

Gene epidermal growth factor receptor (erythroblastic leukemia viral (v-erb-b)

Description: oncogene homolog, avian)

Omim ID: <u>131550</u>, <u>211980</u>

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a transmembrane glycoprotein that

is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this

gene are associated with lung cancer. [provided by RefSeq

Other avian erythroblastic leukemia viral (v-erb-b) oncogene homolog,cell

Designations: growth inhibiting protein 40,cell proliferation-inducing protein

61, epidermal growth factor receptor

Gene Information

Entrez GenelD: 1950

Gene Name: EGF

Gene Alias: HOMG4,URG

Gene epidermal growth factor (beta-urogastrone)

Description:


Omim ID: <u>131530</u>

Gene Ontology: Hyperlink

Gene Summary: Epidermal growth factor has a profound effect on the differentiation of

specific cells in vivo and is a potent mitogenic factor for a variety of cultured cells of both ectodermal and mesodermal origin. The EGF precursor is believed to exist as a membrane-bound molecule which is proteolytically cleaved to generate the 53-amino acid peptide hormone

that stimulates cells to divide. [provided by RefSeq



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/19