

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





TP53 & SMAD2 Protein Protein Interaction Antibody Pair

Catalog #: DI0251 規格:[1 Set]

List All

Specification

Product Description:

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the TP53 protein, and the other against the SMAD2 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




1. Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



3. Hybridize connector oligos



4. Ligation to form a complete DNA circle



5. Rolling circle amplification




6. Add fluorescent probes to reveal interaction

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.

Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between TP53 and SMAD2. HeLa cells were stained with anti-TP53 rabbit purified polyclonal antibody 1:1200 and anti-SMAD2 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied **Product:**

Antibody pair set content:

- 1. TP53 rabbit purified polyclonal antibody (20 ug)
- 2. SMAD2 mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

MSDS:

<u> Download</u>

Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

SMAD2 TP53

Gene Information

Entrez GeneID: 7157

Gene Name: TP53

Gene Alias: FLJ92943,LFS1,TRP53,p53

Gene tumor protein p53

Description:

Omim ID: <u>114480, 114500, 114550, 151623, 161550, 191170, 202300, 260350</u>

Gene Ontology: Hyperlink

Gene Summary: This gene encodes tumor protein p53, which responds to diverse

cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of this gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome. Multiple p53 variants due to alternative promoters and multiple alternative splicing have been found. These variants encode distinct isoforms, which can regulate p53 transcriptional activity. [provided by RefSeq

Other p53

p53 antigen,p53 transformation suppressor,p53 tumor

Designations: suppressor, phosphoprotein p53, transformation-related protein 53

Gene Information

Entrez GeneID: 4087

Gene Name: SMAD2

Gene Alias: JV18,JV18-1,MADH2,MADR2,MGC22139,MGC34440,hMAD-2,hSMAD2

Gene SMAD family member 2

Description:

Page 2 of 3 2016/5/19


Omim ID: 601366

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq

Other **Designations:** MAD, mothers against decapentaplegic homolog 2, Mad protein homolog, Mad, mothers against decapentaplegic homolog 2, Mad-related protein 2,SMAD, mothers against DPP homolog 2,Sma- and Madrelated protein 2, mother against DPP homolog 2

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/19