

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com




## PDGFRB & PIK3R3 Protein Protein Interaction Antibody Pair

Catalog # : DI0262


規格:[1Set]

List All




Quality Control Protein protein interaction immunofluorescence result.

Testing:



	20°C storage immediately after use.
MSDS:	m Download
Publication Ref	erence
novel prognos Liu CH, Chen Cheng HC, Cl	<u>f protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma.</u> TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, hen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 1013 Feb 8. [Epub ahead of print]
Applications	
<i>In situ</i> Proximity	/ Ligation Assay (Cell)
PDGFRB PIK3R	3
Gene Informatio	on
Entrez GenelD:	<u>5159</u>
Gene Name:	PDGFRB
Gene Alias:	CD140B,JTK12,PDGF-R-beta,PDGFR,PDGFR1
Gene Description:	platelet-derived growth factor receptor, beta polypeptide
Omim ID:	<u>131440, 173410</u>
Gene Ontology	: <u>Hyperlink</u>
Gene Summary	This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet- derived growth factor receptor alpha and beta polypeptides. This gene is flanked on chromosome 5 by the genes for granulocyte-macrophage colony-stimulating factor and macrophage-colony stimulating factor receptor; all three genes may be implicated in the 5-q syndrome. A translocation between chromosomes 5 and 12, that fuses this gene to that of the translocation, ETV6, leukemia gene, results in chronic myeloproliferative disorder with eosinophilia. [provided by RefSeq
Other Designations:	beta platelet-derived growth factor receptor,platelet-derived growth factor receptor beta,soluble PDGFRb variant 1
Gene Informatio	on
Entrez GenelD:	8503
Gene Name:	PIK3R3
Gene Alias:	DKFZp686P05226,FLJ41892,p55,p55-GAMMA
Gene Description:	phosphoinositide-3-kinase, regulatory subunit 3 (gamma)
Omim ID:	<u>606076</u>
Gene Ontology	: <u>Hyperlink</u>
Other Designations:	100% homology to SWISS-PROT Q92569,OTTHUMP00000009783,OTTHUMP00000009786,phosphatidyl
	Page 2 of 3



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.