

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



## PDGFRA & PIK3R1 Protein Protein Interaction Antibody Pair

Catalog # : DI0267

規格 : [1 Set]

List All

Specification				Application Image
Product Description:	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the PDGFRA protein, and the other against the PIK3R1 protein for use in <u>in</u> <u>situ</u> Proximity Ligation Assay. See Publication Reference below.			In situ Proximity Ligation Assay (Cell)
	1. Incubate with target	2. Add PLA probes	3. Hybridize connector	
	Primary antibodies	PLUS and MINUS	oligos 6. Add fluorescent probes to reveal interaction	

Reactivity: Human

**Quality Control** Protein protein interaction immunofluorescence result.

Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between PDGFRA and PIK3R1. HeLa cells were stained with anti-PDGFRA rabbit purified polyclonal antibody 1:1200 and anti-PIK3R1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (<u>BlobFinder</u>) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:	Antibody pair set content: 1. PDGFRA rabbit purified polyclonal antibody (20 ug) 2. PIK3R1 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

	20°C storage immediately after use.		
MSDS:	Download		
Publication Ref	erence		
novel prognos Liu CH, Chen Cheng HC, Cl	f protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma. TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, hen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 013 Feb 8. [Epub ahead of print]		
Applications			
In situ Proximity	/ Ligation Assay (Cell)		
PDGFRA PIK3R	1		
Gene Informatio	on		
Entrez GenelD:	5156		
Gene Name:	PDGFRA		
Gene Alias:	CD140A,MGC74795,PDGFR2,Rhe-PDGFRA		
Gene Description:	platelet-derived growth factor receptor, alpha polypeptide		
Omim ID:	<u>173490, 606764, 607685</u>		
Gene Ontology	: <u>Hyperlink</u>		
Gene Summary	This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet- derived growth factor receptor alpha and beta polypeptides. Studies in knockout mice, where homozygosity is lethal, indicate that the alpha form of the platelet-derived growth factor receptor is particularly important for kidney development since mice heterozygous for the receptor exhibit defective kidney phenotypes. [provided by RefSeq		
Other Designations:	FIP1L1/PDGFRA fusion protein,platelet-derived growth factor receptor alpha,rearranged-in-hypereosinophilia-platelet derived growth factor receptor alpha fusion protein		
Gene Informatio	on		
Entrez GenelD:	5295		
Gene Name:	PIK3R1		
Gene Alias:	GRB1,p85,p85-ALPHA		
Gene Description:	phosphoinositide-3-kinase, regulatory subunit 1 (alpha)		
Omim ID:	<u>171833</u>		
Gene Ontology	: <u>Hyperlink</u>		
Gene Summary	Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50		



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.