

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





IGF1R & PIK3R1 Protein Protein Interaction Antibody Pair

Catalog #: DI0269 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the IGF1R protein, and the other against the PIK3R1 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)



 Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



 Hybridize connector oligos



 Ligation to form a complete DNA circle



5. Rolling circle amplification




Add fluorescent probes
to reveal interaction

Reactivity:

Human

Quality Contro

Quality Control Protein protein interaction immunofluorescence result.



Representative image of Proximity Ligation Assay of protein-protein interactions between IGF1R and PlK3R1. HeLa cells were stained with anti-IGF1R rabbit purified polyclonal antibody 1:1200 and anti-PlK3R1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:

Antibody pair set content:

- 1. IGF1R rabbit purified polyclonal antibody (20 ug)
- 2. PIK3R1 mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

MSDS:

<u>m</u>Download

Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

IGF1R PIK3R1

Gene Information

Entrez GeneID: 3480

Gene Name: IGF1R

Gene Alias: CD221,IGFIR,JTK13,MGC142170,MGC142172,MGC18216

Gene insulin-like growth factor 1 receptor

Description:

Omim ID: <u>147370</u>

Gene Ontology: Hyperlink

Gene Summary: This receptor binds insulin-like growth factor with a high affinity. It has

tyrosine kinase activity. The insulin-like growth factor I receptor plays a critical role in transformation events. Cleavage of the precursor generates alpha and beta subunits. It is highly overexpressed in most malignant tissues where it functions as an anti-apoptotic agent by

enhancing cell survival. [provided by RefSeq

Other soluble IGF1R variant 1, soluble IGF1R variant 2

Designations:

Gene Information

Entrez GenelD: 5295

Gene Name: PIK3R1

Gene Alias: GRB1,p85,p85-ALPHA

Gene phosphoinositide-3-kinase, regulatory subunit 1 (alpha)

Description:

Omim ID: <u>171833</u>

Gene Ontology: Hyperlink

Gene Summary: Phosphatidylinositol 3-kinase phosphorylates the inositol ring of

phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50

kD. This gene encodes the 85 kD regulatory subunit.

Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance. Alternative splicing of this gene results in three transcript variants encoding different isoforms. [provided by RefSeq


Page 2 of 3 2016/5/19

Other Designations:

phosphatidylinositol 3-kinase, regulatory subunit, polypeptide 1 (p85 alpha),phosphatidylinositol 3-kinase, regulatory, 1,phosphatidylinositol 3-kinase-associated p-85 alpha,phosphoinositide-3-kinase, regulatory

subunit 1 (p85 alpha),phosphoinositide-3-ki

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/19