

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





EGFR & FOS Protein Protein Interaction Antibody Pair

Catalog #: DI0275 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the EGFR protein, and the other against the FOS protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image

In situ Proximity Ligation Assay (Cell)




1. Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS




3. Hybridize connector oligos



4. Ligation to form a complete DNA circle



5. Rolling circle amplification




6. Add fluorescent probes to reveal interaction

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.





Representative image of Proximity Ligation Assay of protein-protein interactions between EGFR and FOS. HeLa cells were stained with anti-EGFR rabbit purified polyclonal antibody 1:1200 and anti-FOS mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied **Product:**

Antibody pair set content:

- 1. EGFR rabbit purified polyclonal antibody (20 ug)
- 2. FOS mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

20°C storage immediately after use.

MSDS:

Download

Proteomics. 2013 Feb 8. [Epub ahead of print]

Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
 Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell

Applications

In situ Proximity Ligation Assay (Cell)

EGFR FOS

Gene Information

Entrez GeneID: 1956

Gene Name: EGFR

Gene Alias: ERBB,ERBB1,HER1,PIG61,mENA

Gene epidermal growth factor receptor (erythroblastic leukemia viral (v-erb-b)

Description: oncogene homolog, avian)

Omim ID: <u>131550</u>, <u>211980</u>

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a transmembrane glycoprotein that

is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this

gene are associated with lung cancer. [provided by RefSeq

Other avian erythroblastic leukemia viral (v-erb-b) oncogene homolog,cell

Designations: growth inhibiting protein 40,cell proliferation-inducing protein

61, epidermal growth factor receptor

Gene Information

Entrez GenelD: 2353

Gene Name: FOS

Gene Alias: AP-1,C-FOS

Gene v-fos FBJ murine osteosarcoma viral oncogene homolog

Description:

Omim ID: <u>164810</u>

Gene Ontology: Hyperlink

Gene Summary: The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and

FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with

apoptotic cell death. [provided by RefSeq

Other FBJ murine osteosarcoma viral (v-fos) oncogene homolog (oncogene FOS), activator protein 1, cellular oncogene c-fos

Interactome

CDH1

CTNNB1

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/19