

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



TRAF2 & MAPK9 Protein Protein Interaction Antibody Pair

Catalog # : DI0282

規格:[1Set]


List All

Specification				Application Image
Product Description:	This protein protein inte antibodies to detect the TRAF2 protein, and the <u>situ Proximity Ligation A</u>	protein-protein interaction other against the MAPK	on, one against the ⑶ protein for use in <u>in</u>	In situ Proximity Ligation Assay (Cell)
		A A		
	 Incubate with target primary antibodies 	2. Add PLA probes PLUS and MINUS	 Hybridize connector oligos 	
	4. Ligation to form a	5. Rolling circle	6. Add fluorescent probes	
	complete DNA circle	amplification	to reveal interaction	

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.



	20°C storage immediately after use.		
MSDS:	ma Download		
Publication Ref	erence		
novel progno Liu CH, Chen Cheng HC, C	of protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma. TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, hen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 2013 Feb 8. [Epub ahead of print]		
Applications			
In situ Proximit	y Ligation Assay (Cell)		
MAPK9 TRAF2			
Gene Informati	on		
Entrez GenelD:	7186		
Gene Name:	TRAF2		
Gene Alias:	MGC:45012,TRAP,TRAP3		
Gene Description:	TNF receptor-associated factor 2		
Omim ID:	<u>601895</u>		
Gene Ontology	: <u>Hyperlink</u>		
Gene Summary	The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can unbiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been found for this gene, but the biological validity of only one transcript has been determined. [provided by RefSeq		
Other Designations:	OTTHUMP00000022625,OTTHUMP00000064745,tumor necrosis factor type 2 receptor associated protein 3		
Gene Informati	on		
Entrez GenelD:	: <u>5601</u>		
Gene Name:	MAPK9		
Gene Alias:	JNK- 55,JNK2,JNK2A,JNK2ALPHA,JNK2B,JNK2BETA,PRKM9,SAPK,p54a,p54 aSAPK		
Gene Description:	mitogen-activated protein kinase 9		
	Page 2 of 3		


Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase targets specific transcription factors, and thus mediates immediate-early gene expression in response to various cell stimuli. It is most closely related to MAPK8, both of which are involved in UV radiation induced apoptosis, thought to be related to the cytochrome cmediated cell death pathway. This gene and MAPK8 are also known as c-Jun N-terminal kinases. This kinase blocks the ubiquitination of tumor suppressor p53, and thus it increases the stability of p53 in nonstressed cells. Studies of this gene's mouse counterpart suggest a key role in Tcell differentiation. Several alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq

Other Designations:

Jun kinase,MAP kinase 9,c-Jun N-terminal kinase 2,c-Jun kinase 2,mitogen-activated protein kinase 9 isoform JNK2 alpha2,stressactivated protein kinase JNK2

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.