

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





XIAP & CASP9 Protein Protein Interaction Antibody Pair

Catalog #: DI0293 規格:[1 Set]

List All

Specification

Product Description:

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the XIAP protein, and the other against the CASP9 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image

In situ Proximity Ligation Assay (Cell)




1. Incubate with target primary antibodies




2. Add PLA probes PLUS and MINUS




3. Hybridize connector oligos



4. Ligation to form a complete DNA circle



5. Rolling circle amplification




6. Add fluorescent probes to reveal interaction

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.

Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between XIAP and CASP9. HeLa cells were stained with anti-XIAP rabbit purified polyclonal antibody 1:1200 and anti-CASP9 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied **Product:**

Antibody pair set content:

- 1. XIAP rabbit purified polyclonal antibody (20 ug)
- 2. CASP9 mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

<u>m</u>Download

Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

XIAP CASP9

Gene Information

Entrez GeneID: 331

Gene Name: XIAP

Gene Alias: API3,BIRC4,ILP1,MIHA,XLP2

Gene X-linked inhibitor of apoptosis

Description:

Omim ID: 300079, 300635

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a member of a family of proteins

which inhibit apoptosis through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2. This protein inhibits apoptosis induced by menadione, a potent inducer of free radicals, and ICE. It also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. [provided by RefSeq

Other Designations:

OTTHUMP00000023975,OTTHUMP00000196392,apoptosis inhibitor

3,baculoviral IAP repeat-containing 4,baculoviral IAP repeat-containing

protein 4

Gene Information

Entrez GeneID: 842

Gene Name: CASP9

Gene Alias: APAF-3,APAF3,CASPASE-9c,ICE-LAP6,MCH6

Gene caspase 9, apoptosis-related cysteine peptidase

Description:

Omim ID: 602234

Gene Ontology: Hyperlink

Gene Summary: This gene encodes a member of the cysteine-aspartic acid protease

(caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein is processed by caspase APAF1; this step is thought to be one of the earliest in the caspase activation

cascade. Alternative splicing results in two transcript variants which encode different isoforms. [provided by RefSeq


Other Designations:

ICE-like apoptotic protease

ns: 6,OTTHUMP0000002322,OTTHUMP0000002323,OTTHUMP0000004 4594,apoptotic protease MCH-6,apoptotic protease activating factor

3,caspase 9,caspase 9, apoptosis-related cysteine protease

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/19