



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## AKT1 & XIAP Protein Protein Interaction Antibody Pair


Catalog # : DI0304

規格 : [ 1 Set ]

List All


### Specification

**Product Description:** This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the AKT1 protein, and the other against the XIAP protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.



**Reactivity:** Human

**Quality Control Testing:** Protein protein interaction immunofluorescence result.




Representative image of Proximity Ligation Analysis of protein-protein interactions between AKT1 and XIAP. HeLa cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-XIAP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware ([BlobFinder](#)) download from The Centre for Image Analysis at Uppsala University.

**Supplied Product:** Antibody pair set content:  
 1. AKT1 rabbit purified polyclonal antibody (20 ug)  
 2. XIAP mouse monoclonal antibody (40 ug)  
 \*Reagents are sufficient for at least 30-50 assays using recommended protocols.

**Storage Instruction:** Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -


### Application Image

*In situ* Proximity Ligation Assay (Cell)




[enlarge](#)

*In situ* Proximity Ligation Assay (Cell)




[enlarge](#)

*In situ* Proximity Ligation Assay (Cell)



[enlarge](#)

*In situ* Proximity Ligation Assay (Cell)



[enlarge](#)

20°C storage immediately after use.

MSDS:

 [Download](#)

---

## Publication Reference


---

1. [An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.](#)  
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]
- 

## Applications

---


### *In situ* Proximity Ligation Assay (Cell)



Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and XIAP. PC-3 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-XIAP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

---


### *In situ* Proximity Ligation Assay (Cell)



Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and XIAP. A-549 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-XIAP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

---


### *In situ* Proximity Ligation Assay (Cell)



Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and XIAP. HT-29 cells were stained with anti-AKT1 rabbit purified polyclonal antibody

1:100 and anti-XIAP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

#### ***In situ* Proximity Ligation Assay (Cell)**



Confocal microscopy image of Proximity Ligation Assay of protein-protein interactions between AKT1 and XIAP. HT-29 cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:100 and anti-XIAP mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).

[AKT1](#) [XIAP](#)

#### **Gene Information**

**Entrez GeneID:** [207](#)

**Gene Name:** AKT1

**Gene Alias:** AKT,MGC99656,PKB,PKB-ALPHA,PRKBA,RAC,RAC-ALPHA

**Gene Description:** v-akt murine thymoma viral oncogene homolog 1

**Omim ID:** [164730](#), [181500](#)

**Gene Ontology:** [Hyperlink](#)

**Gene Summary:** The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq]

**Other Designations:** RAC-alpha serine/threonine-protein kinase,murine thymoma viral (v-akt) oncogene homolog-1,protein kinase B,rac protein kinase alpha

#### **Gene Information**

**Entrez GeneID:** [331](#)

**Gene Name:** XIAP

**Gene Alias:** API3,BIRC4,ILP1,MIHA,XLP2

**Gene Description:** X-linked inhibitor of apoptosis


Omim ID: 300079, 300635

Gene Ontology: [Hyperlink](#)

**Gene Summary:** The protein encoded by this gene is a member of a family of proteins which inhibit apoptosis through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2. This protein inhibits apoptosis induced by menadione, a potent inducer of free radicals, and ICE. It also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. [provided by RefSeq]

**Other Designations:** OTTHUMP00000023975, OTTHUMP00000196392, apoptosis inhibitor 3, baculoviral IAP repeat-containing 4, baculoviral IAP repeat-containing protein 4

### Interactome



[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.