

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





MAPK14 & MAPKAPK3 Protein Protein Interaction Antibody Pair

Catalog #: DI0344 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the MAPK14 protein, and the other against the MAPKAPK3 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




 Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



 Hybridize connector oligos



 Ligation to form a complete DNA circle



Rolling circle amplification




Add fluorescent probes
to reveal interaction

Reactivity:

Human

Quality Contro

Quality Control Protein protein interaction immunofluorescence result.



Representative image of Proximity Ligation Assay of protein-protein interactions between MAPK14 and MAPKAPK3. HeLa cells were stained with anti-MAPK14 rabbit purified polyclonal antibody 1:1200 and anti-MAPKAPK3 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:

Antibody pair set content:

- 1. MAPK14 rabbit purified polyclonal antibody (20 ug)
- 2. MAPKAPK3 mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

20°C storage immediately after use.

MSDS:

<u>m</u>Download

Publication Reference

1. An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma. Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

MAPK14 MAPKAPK3

Gene Information

Entrez GeneID: 1432

Gene Name: MAPK14

Gene Alias: CSBP1,CSBP2,CSPB1,EXIP,Mxi2,PRKM14,PRKM15,RK,SAPK2A,p38,p

38ALPHA

Gene mitogen-activated protein kinase 14

Description:

Omim ID: 600289

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a member of the MAP kinase

family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct

Other

Csaids binding protein, MAP kinase Mxi2, MAX-interacting protein 2,cytokine suppressive anti-inflammatory drug binding protein,p38 MAP **Designations:**

isoforms have been reported. [provided by RefSeq

kinase,p38 mitogen activated protein kinase,p38alpha Exip,stress-

activated protein kinase 2A

Gene Information

Entrez GeneID: 7867

Gene Name: MAPKAPK3

Gene Alias: 3PK,MAPKAP3

mitogen-activated protein kinase-activated protein kinase 3 Gene

Description:

602130 Omim ID:


> Page 2 of 3 2016/5/19

Gene Ontology: Hyperlink

Gene Summary: This gene encodes a member of the Ser/Thr protein kinase family. This kinase functions as a mitogen-activated protein kinase (MAP kinase)activated protein kinase. MAP kinases are also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This kinase was shown to be activated by growth inducers and stress stimulation of cells. In vitro studies demonstrated that ERK, p38 MAP kinase and Jun N-terminal kinase were all able to phosphorylate and activate this kinase, which suggested the role of this kinase as an integrative element of signaling in both mitogen and stress responses. This kinase was reported to interact with, phosphorylate and repress the activity of E47, which is a basic helix-loop-helix transcription factor known to be involved in the regulation of tissue-specific gene expression and cell differentiation. [provided by RefSeq

Other **Designations:** MAPKAP kinase 3

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

2016/5/19 Page 3 of 3