

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



TP53 & PPP2R5C Protein Protein Interaction Antibody Pair

Catalog # : DI0378

規格:[1Set]

List All

Specification				Application Image
Product Description:	This protein protein inte antibodies to detect the TP53 protein, and the o <u>situ</u> Proximity Ligation A	protein-protein interact ther against the PPP2F	ion, one against the R5C protein for use in <u>in</u>	<i>In situ</i> Proximity Ligation Assay (Cell)
	1. Incubate with target primary antibodies	2. Add PLA probes PLUS and MINUS	3. Hybridize connector oligos	
	4. Ligation to form a complete DNA circle	5. Rolling circle amplification	6. Add fluorescent probes to reveal interaction	

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.


Testina:

looning.	Representative image of Proximity Ligation Assay of protein-protein interactions between TP53 and PPP2R5C. HeLa cells were stained with anti-TP53 rabbit purified polyclonal antibody 1:1200 and anti-PPP2R5C mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.
Supplied Product:	Antibody pair set content: 1. TP53 rabbit purified polyclonal antibody (20 ug) 2. PPP2R5C mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

	20°C storage immediately after use.
MSDS:	Download
Publication Re	ference
novel progno Liu CH, Cher Cheng HC, C Proteomics.	of protein-protein interactions in cross-talk pathways reveals CRKL as a ostic marker in hepatocellular carcinoma. In TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 2013 Feb 8. [Epub ahead of print]
Applications In situ Proximit	y Ligation Assay (Cell)
PPP2R5C TP53	<u>1</u>
Gene Informati	ion
Entrez GenelD	: <u>7157</u>
Gene Name:	TP53
Gene Alias:	FLJ92943,LFS1,TRP53,p53
Gene Description:	tumor protein p53
Omim ID:	<u>114480, 114500, 114550, 151623, 161550, 191170, 202300, 260350</u>
Gene Ontology	r: <u>Hyperlink</u>
Gene Summary	<i>r</i> : This gene encodes tumor protein p53, which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of this gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome. Multiple p53 variants due to alternative promoters and multiple alternative splicing have been found. These variants encode distinct isoforms, which can regulate p53 transcriptional activity. [provided by RefSeq
Other Designations:	p53 antigen,p53 transformation suppressor,p53 tumor suppressor,phosphoprotein p53,transformation-related protein 53
Gene Informat	on
Entrez GenelD	: 5527
Gene Name:	PPP2R5C
Gene Alias:	B56G,MGC23064,PR61G
Gene Description:	protein phosphatase 2, regulatory subunit B', gamma isoform

Omim ID:	<u>601645</u>		
Gene Ontology	Hyperlink		
Gene Summary	The product of this gene belongs to the phosphatase 2A regulatory subunit B family. Protein phosphatase 2A is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The B regulatory subunit might modulate substrate selectivity and catalytic activity. This gene encodes a gamma isoform of the regulatory subunit B56 subfamily. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq		
DtherB' alpha regulatory subunit,PP2A, B subunit, B' gamma isoform,PDesignations:subunit, B56 gamma isoform,PP2A, B subunit, PR61 gammaisoform,PP2A, B subunit, R5 gamma isoform,Serine/threonine prophosphatase 2A, 56 kDa regulatory subunit, gamma isoform,gam			

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.