

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





AKT1 & PPM1A Protein Protein Interaction Antibody Pair

Catalog #: DI0384 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the AKT1 protein, and the other against the PPM1A protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




1. Incubate with target primary antibodies




2. Add PLA probes PLUS and MINUS



3. Hybridize connector oligos



4. Ligation to form a complete DNA circle



5. Rolling circle amplification




6. Add fluorescent probes to reveal interaction

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.





Representative image of Proximity Ligation Assay of protein-protein interactions between AKT1 and PPM1A. HeLa cells were stained with anti-AKT1 rabbit purified polyclonal antibody 1:1200 and anti-PPM1A mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied **Product:**

Antibody pair set content:

- 1. AKT1 rabbit purified polyclonal antibody (20 ug)
- 2. PPM1A mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -



Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell

Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

AKT1 PPM1A

Gene Information

Entrez GeneID: 207

Gene Name: AKT1

Gene Alias: AKT,MGC99656,PKB,PKB-ALPHA,PRKBA,RAC,RAC-ALPHA

Gene v-akt murine thymoma viral oncogene homolog 1

Description:

Omim ID: <u>164730</u>, <u>181500</u>

Gene Ontology: Hyperlink

Gene Summary: The serine-threonine protein kinase encoded by the AKT1 gene is

catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Multiple alternatively spliced transcript variants

have been found for this gene. [provided by RefSeq

Other RAC-alpha serine/threonine-protein kinase,murine thymoma viral (v-akt)

Designations: oncogene homolog-1, protein kinase B, rac protein kinase alpha

Gene Information

Entrez GeneID: 5494

Gene Name: PPM1A

Gene Alias: FLJ42306,MGC9201,PP2C-ALPHA,PP2CA

Gene protein phosphatase 1A (formerly 2C), magnesium-dependent, alpha

Description: isoform

Omim ID: 606108

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a member of the PP2C family of

Ser/Thr protein phosphatases. PP2C family members are known to be


Page 2 of 3 2016/5/19

negative regulators of cell stress response pathways. This phosphatase dephosphorylates, and negatively regulates the activities of, MAP kinases and MAP kinases kinases. It has been shown to inhibit the activation of p38 and JNK kinase cascades induced by environmental stresses. This phosphatase can also dephosphorylate cyclin-dependent kinases, and thus may be involved in cell cycle control. Overexpression of this phosphatase is reported to activate the expression of the tumor suppressor gene TP53/p53, which leads to G2/M cell cycle arrest and apoptosis. Three alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq

Other Designations:

protein phosphatase 1A, protein phosphatase 2C alpha isoform

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/19