

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com




### CDC6 & MCM6 Protein Protein Interaction Antibody Pair


Catalog #: D10399

規格:[1Set]

List All



Quality Control Protein protein interaction immunofluorescence result.



#### 20°C storage immediately after use.


MSDS: million

**Publication Reference** 

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
 Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

#### **Applications**

In situ Proximity Ligation Assay (Cell)



Representative image of Proximity Ligation Assay of protein-protein interactions between CDC6 and MCM6. Mahlavu cells were stained with anti-CDC6 rabbit purified polyclonal antibody 1:1200 and anti-MCM6 mouse purified polyclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex, and nuclei were counterstained with DAPI (blue).


#### CDC6 MCM6

#### **Gene Information**

Entrez GenelD:	<u>990</u>
Gene Name:	CDC6
Gene Alias:	CDC18L,HsCDC18,HsCDC6
Gene Description:	cell division cycle 6 homolog (S. cerevisiae)
Omim ID:	602627
Gene Ontology:	Hyperlink
Gene Summary:	The protein encoded by this gene is highly similar to Saccharomyces cerevisiae Cdc6, a protein essential for the initiation of DNA replication. This protein functions as a regulator at the early steps of DNA replication. It localizes in cell nucleus during cell cyle G1, but translocates to the cytoplasm at the start of S phase. The subcellular translocation of this protein during cell cyle is regulated through its phosphorylation by Cdks. Transcription of this protein was reported to be regulated in response to mitogenic signals through transcriptional control mechanism involving E2F proteins. [provided by RefSeq
Other Designations:	CDC18 (cell division cycle 18, S.pombe, homolog)-like,CDC6 cell division cycle 6 homolog,CDC6-related protein,cell division cycle 6 protein
Gene Informatio	ภ

Entrez GenelD:	<u>4175</u>
Gene Name:	MCM6
Gene Alias:	MCG40308, Mis5, P105MCM
Gene Description:	minichromosome maintenance complex component 6
Omim ID:	<u>223100, 601806</u>
Gene Ontology	: <u>Hyperlink</u>
Gene Summary	: The protein encoded by this gene is one of the highly conserved mini- chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre- replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 7 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. The phosphorylation of the complex by CDC2 kinase reduces the helicase activity, suggesting a role in the regulation of DNA replication. [provided by RefSeq
Other Designations:	DNA replication licensing factor MCM6,MCM6 minichromosome maintenance deficient 6 (MIS5 homolog, S. pombe),MIS5 homolog,minichromosome maintenance deficient (mis5, S. pombe) 6,minichromosome maintenance deficient 6 homolog

#### Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.