

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





PDGFRB & PTEN Protein Protein Interaction Antibody Pair

Catalog #: DI0425 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the PDGFRB protein, and the other against the PTEN protein for use in <u>in situ Proximity Ligation Assay</u>. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




 Incubate with target primary antibodies




2. Add PLA probes PLUS and MINUS



 Hybridize connector oligos



 Ligation to form a complete DNA circle



Rolling circle amplification




Add fluorescent probes
to reveal interaction

Reactivity:

Human

Quality Contro Testing:

Quality Control Protein protein interaction immunofluorescence result.



Representative image of Proximity Ligation Assay of protein-protein interactions between PDGFRB and PTEN. HeLa cells were stained with anti-PDGFRB rabbit purified polyclonal antibody 1:1200 and anti-PTEN mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:

Antibody pair set content:

- 1. PDGFRB rabbit purified polyclonal antibody (20 ug)
- 2. PTEN mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

MSDS:

<u> Download</u>

Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

PDGFRB PTEN

Gene Information

Entrez GeneID: 5159

Gene Name: PDGFRB

Gene Alias: CD140B,JTK12,PDGF-R-beta,PDGFR,PDGFR1

Gene platelet-derived growth factor receptor, beta polypeptide

Description:

Omim ID: <u>131440</u>, <u>173410</u>

Gene Ontology: Hyperlink

Gene Summary: This gene encodes a cell surface tyrosine kinase receptor for members

of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. This gene is flanked on chromosome 5 by the genes for granulocyte-macrophage colony-stimulating factor and macrophage-colony stimulating factor receptor; all three genes may be implicated in the 5-q syndrome. A translocation between chromosomes 5 and 12, that fuses this gene to that of the translocation, ETV6, leukemia gene, results in chronic myeloproliferative disorder with eosinophilia. [provided by RefSeq

Other beta platelet-derived growth factor receptor, platelet-derived growth

Designations: factor receptor beta, soluble PDGFRb variant 1

Gene Information

Entrez GeneID: 5728

Gene Name: PTEN

Gene Alias: 10q23del,BZS,MGC11227,MHAM,MMAC1,PTEN1,TEP1

Gene phosphatase and tensin homolog

Description:

Omim ID: <u>137800</u>, <u>153480</u>, <u>158350</u>, <u>174900</u>, <u>176807</u>, <u>176920</u>, <u>188470</u>, <u>276950</u>,

601728, 605309, 607174

Gene Ontology: Hyperlink

Gene Summary: This gene was identified as a tumor suppressor that is mutated in a


Page 2 of 3 2016/5/20

large number of cancers at high frequency. The protein encoded this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. [provided by RefSeq

Other Designations:

MMAC1 phosphatase and tensin homolog deleted on chromosome 10,OTTHUMP0000020032,mutated in multiple advanced cancers 1

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/20