

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



PTK2 & STAT3 Protein Protein Interaction Antibody Pair

Catalog # : DI0435


規格 : [1 Set]

List All

Specification				Application Image
Product Description:	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the PTK2 protein, and the other against the STAT3 protein for use in <u>in situ</u> <u>Proximity Ligation Assay</u> . <u>See Publication Reference below</u> .			In situ Proximity Ligation Assay (Cell)
		A A	A A A A A A A A A A A A A A A A A A A	
	1. Incubate with target primary antibodies	2. Add PLA probes PLUS and MINUS	 Hybridize connector oligos 	
	4. Ligation to form a	5. Rolling circle	6. Add fluorescent probes	
	4. Ligation to form a complete DNA circle	5. Rolling circle amplification	6. Add fluorescent probes to reveal interaction	

Reactivity: Human

Quality Control Protein protein interaction immunofluorescence result.




	20°C storage immediately after use.			
MSDS:	Download			
Publication Ref	erence			
novel progno Liu CH, Chen Cheng HC, C	f protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma. TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, hen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 2013 Feb 8. [Epub ahead of print]			
Applications				
<i>In situ</i> Proximity	y Ligation Assay (Cell)			
PTK2 STAT3				
Gene Informati	on			
Entrez GenelD:	<u>5747</u>			
Gene Name:	РТК2			
Gene Alias:	FADK,FAK,FAK1,pp125FAK			
Gene Description:	PTK2 protein tyrosine kinase 2			
Omim ID:	600758			
Gene Ontology	: <u>Hyperlink</u>			
Gene Summary	This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. At least four transcript variants encoding four different isoforms have been found for this gene, but the full-length natures of only two of them have been determined. [provided by RefSeq			
Other Designations:	focal adhesion kinase 1			
Gene Informati	on			
Entrez GenelD:	<u>6774</u>			
Gene Name:	STAT3			
Gene Alias:	APRF,FLJ20882,HIES,MGC16063			
Gene Description:	signal transducer and activator of transcription 3 (acute-phase response factor)			
Omim ID:	102582			
Gene Ontology	: <u>Hyperlink</u>			
Gene Summary	: The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and			

then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated
through phosphorylation in response to various cytokines and growth
factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli,
and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and
regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Three alternatively spliced transcript variants encoding
distinct isoforms have been described. [provided by RefSeq

OtherDNA-binding protein APRF,acute-phase response factor,signalDesignations:transducer and activator of transcription 3

Interactome



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.