

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





## FAS & RHOA Protein Protein Interaction Antibody Pair

Catalog # : DI0454 規格:[1 Set]

#### List All

## **Specification**

## **Product Description:**

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the FAS protein, and the other against the RHOA protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

## **Application Image**


In situ Proximity Ligation Assay (Cell)




1. Incubate with target primary antibodies




2. Add PLA probes PLUS and MINUS



3. Hybridize connector oligos



4. Ligation to form a complete DNA circle



5. Rolling circle amplification




6. Add fluorescent probes to reveal interaction

Reactivity:

Human

Quality Control Protein protein interaction immunofluorescence result.

Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between FAS and RHOA. HeLa cells were stained with anti-FAS rabbit purified polyclonal antibody 1:1200 and anti-RHOA mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

## Supplied **Product:**

Antibody pair set content:

- 1. FAS rabbit purified polyclonal antibody (20 ug)
- 2. RHOA mouse monoclonal antibody (40 ug)

\*Reagents are sufficient for at least 30-50 assays using recommended protocols.

### **Storage** Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

20°C storage immediately after use.

MSDS:

<u>m</u>Download

#### **Publication Reference**

1. An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma. Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

#### **Applications**

In situ Proximity Ligation Assay (Cell)

FAS RHOA

#### **Gene Information**

Entrez GeneID: 355

Gene Name: FAS

Gene Alias: ALPS1A,APO-1,APT1,CD95,FAS1,FASTM,TNFRSF6

Fas (TNF receptor superfamily, member 6) Gene

**Description:** 

Omim ID: <u>134637</u>, <u>601859</u>

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains a death domain. It has been shown to play a central role in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. The interaction of this receptor with its ligand allows the formation of a death-inducing signaling complex that includes Fas-associated death domain protein (FADD), caspase 8, and caspase 10. The autoproteolytic processing of the caspases in the complex triggers a downstream caspase cascade, and leads to apoptosis. This receptor has been also shown to activate NF-kappaB, MAPK3/ERK1, and MAPK8/JNK, and is found to be involved in transducing the proliferating signals in normal diploid fibroblast and T cells. At least eight alternatively spliced transcript variants have been described, some of which are candidates for nonsense-mediated decay (NMD). The isoforms lacking the transmembrane domain may negatively regulate the apoptosis mediated by the full length isoform. [provided by RefSeq

Other Designations: APO-1 cell surface antigen, CD95 antigen, Fas AMA, Fas antigen,OTTHUMP00000020045,OTTHUMP00000020046,OTTHUMP00 000020051,OTTHUMP00000059646,apoptosis antigen 1,tumor necrosis factor receptor superfamily member 6, tumor necrosis factor receptor superfamily, mem

## Gene Information

Entrez GenelD: 387

Gene Name: RHOA

Gene Alias: ARH12,ARHA,RHO12,RHOH12

Gene ras homolog gene family, member A

**Description:** 

Page 2 of 3 2016/5/20 Omim ID: <u>165390</u> Gene Ontology: Hyperlink Aplysia ras-related homolog 12,oncogene RHO H12,small GTP binding Other Designations: protein RhoA Interactome PTPN13 **EGFR** CASP8 MET CTNNB1 FYN FAS RUVBL2 EZR PTPN6

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/20