

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





RALBP1 & TRAF2 Protein Protein Interaction Antibody Pair

Catalog #: DI0471 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the RALBP1 protein, and the other against the TRAF2 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




 Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



 Hybridize connector oligos



 Ligation to form a complete DNA circle

Human



5. Rolling circle amplification




Add fluorescent probes
to reveal interaction

Reactivity:

Quality Contro

Quality Control Protein protein interaction immunofluorescence result.



Representative image of Proximity Ligation Assay of protein-protein interactions between RALBP1 and TRAF2. HeLa cells were stained with anti-RALBP1 rabbit purified polyclonal antibody 1:1200 and anti-TRAF2 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:

Antibody pair set content:

- 1. RALBP1 rabbit purified polyclonal antibody (20 ug)
- 2. TRAF2 mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

2016/5/20

MSDS:

Publication Reference

 An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.
Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

TRAF2 RALBP1

Gene Information

Entrez GeneID: 10928

Gene Name: RALBP1

Gene Alias: RIP1,RLIP1,RLIP76

Gene ralA binding protein 1

Description:

Omim ID: <u>605801</u>

Gene Ontology: Hyperlink

Other OTTHUMP00000162351,ral-interacting protein 1,ralA-binding protein 1

Designations:

Gene Information

Entrez GeneID: 7186

Gene Name: TRAF2

Gene Alias: MGC:45012,TRAP,TRAP3

Gene TNF receptor-associated factor 2

Description:

Omim ID: <u>601895</u>

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene is a member of the TNF receptor

associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can unbiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been found for this gene,

but the biological validity of only one transcript has been determined. [provided by RefSeq Other OTTHUMP00000022625,OTTHUMP0000064745,tumor necrosis factor type 2 receptor associated protein 3 Designations: Interactome BIRC3 TUBB CASP8 TRPC4AP HSPA1A TUBA3C MAP3K8 MAP3K5_{PSEN1} BCL₁₀ CALM3 CLTC TUBB4 TANK HSP90AA1 HSPA5 TNFRSF1A EEF1A2 CAV1 NR2C2 TAOK3 ERN1 CALM₁ HSP90AB1 TRAF2 TNFAIR3 LN/ HSPA9 TRAF6 RIPK3 KBKE MAP3K14 MAP3K7IP2 MAP3K1 TRAF3(P2 ABLIMA TNFRSF1B PRIM1 TRADD TRAF1 IKBKB NFKBIA MAP3K7 RIPK1 NFKB2 RIPK2 ARRB1 GRB₂ IKBKG ARRB2MAP3K3 RELB RELA NFKBIB NFKB1

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/20