

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





CRK & FLNA Protein Protein Interaction Antibody Pair

Catalog #: DI0507 規格:[1 Set]

List All

Specification

Product Description:


This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the CRK protein, and the other against the FLNA protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)




 Incubate with target primary antibodies



2. Add PLA probes PLUS and MINUS



 Hybridize connector oligos



 Ligation to form a complete DNA circle



 Rolling circle amplification




Add fluorescent probes
 to reveal interaction

Reactivity:

Human

Quality Control Testing:

Quality Control Protein protein interaction immunofluorescence result.



Representative image of Proximity Ligation Assay of protein-protein interactions between CRK and FLNA. HeLa cells were stained with anti-CRK rabbit purified polyclonal antibody 1:1200 and anti-FLNA mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:

Antibody pair set content:

- 1. CRK rabbit purified polyclonal antibody (20 ug)
- 2. FLNA mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

MSDS:

<u> Download</u>

Publication Reference

An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.

Liu CH, Chon TC, Chau GV, Ion VH, Chon CH, Heu CN, Lin KT, Iuang VI, Liu PJ, Chon CH, Heu CN, Lin KT, Iuang VI, Liu PJ, Chon CH, Heu CN, Lin KT, Iuang VI, Liu PJ, Chon CH, Heu CN, Lin KT, Iuang VI, Liu PJ, Chon CH, Heu CN, Lin KT, Iuang VI, Liu PJ, Chon CH, Heu CN, Lin KT, Iuang VI, Liu PJ, Chon CH, Heu CN, Lin KT, Iuang VI, Liu PJ, Chon CH, Heu CN, Liu PJ, Chon CH, Chon

Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

CRK FLNA

Gene Information

Entrez GeneID: 1398

Gene Name: CRK

Gene Alias: CRKII

Gene v-crk sarcoma virus CT10 oncogene homolog (avian)

Description:

Omim ID: <u>164762</u>

Gene Ontology: Hyperlink

Gene Summary: This gene encodes a member of an adapter protein family that binds to

several tyrosine-phosphorylated proteins. The product of this gene has several SH2 and SH3 domains (src-homology domains) and is involved in several signaling pathways, recruiting cytoplasmic proteins in the vicinity of tyrosine kinase through SH2-phosphotyrosine interaction. The N-terminal SH2 domain of this protein functions as a positive regulator of transformation whereas the C-terminal SH3 domain functions as a negative regulator of transformation. Two alternative transcripts encoding different isoforms with distinct biological activity have been

described. [provided by RefSeq

Other avian sarcoma virus CT10 (v-crk) oncogene homolog,v-crk avian sarcoma virus CT10 oncogene homolog,v-crk sarcoma virus CT10

oncogene homolog

Gene Information

Entrez GeneID: 2316

Gene Name: FLNA

Gene Alias: ABP-

280,ABPX,DKFZp434P031,FLN,FLN1,FMD,MNS,NHBP,OPD,OPD1,OPD

2

Gene filamin A, alpha (actin binding protein 280)

Description:

Omim ID: 300017, 300049, 300537, 304120, 309350, 311300

Gene Ontology: <u>Hyperlink</u>


Gene Summary: The protein encoded by this gene is an actin-binding protein that

Page 2 of 4 2016/5/20


crosslinks actin filaments and links actin filaments to membrane glycoproteins. The encoded protein is involved in remodeling the cytoskeleton to effect changes in cell shape and migration. This protein interacts with integrins, transmembrane receptor complexes, and second messengers. Defects in this gene are a cause of several syndromes, including periventricular nodular heterotopias (PVNH1, PVNH4), otopalatodigital syndromes (OPD1, OPD2), frontometaphyseal dysplasia (FMD), Melnick-Needles syndrome (MNS), and X-linked congenital idiopathic intestinal pseudoobstruction (CIIPX). Two transcript variants encoding different isoforms have been found for this gene

Other **Designations:** OTTHUMP00000024320, actin-binding protein 280, filamin 1, filamin A,

Interactome 1



Interactome 2



Page 4 of 4 2016/5/20