

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



#### MAP2K6 & TAOK2 Protein Protein Interaction Antibody Pair

#### Catalog # : DI0515

規格:[1Set]

List All


Specification				Application Image
Product Description:	This protein protein inte antibodies to detect the MAP2K6 protein, and th <u>situ Proximity Ligation A</u>	protein-protein interact e other against the TAC	ion, one against the DK2 protein for use in <u>in</u>	<i>In situ</i> Proximity Ligation Assay (Cell)
	1. Incubate with target primary antibodies	2. Add PLA probes PLUS and MINUS	3. Hybridize connector oligos	
	4. Ligation to form a complete DNA circle	5. Rolling circle amplification	6. Add fluorescent probes to reveal interaction	

**Reactivity:** 

Human

Quality Control Protein protein interaction immunofluorescence result.

Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between MAP2K6 and TAOK2. HeLa cells were stained with anti-MAP2K6 rabbit purified polyclonal antibody 1:1200 and anti-TAOK2 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:	Antibody pair set content: 1. MAP2K6 rabbit purified polyclonal antibody (20 ug) 2. TAOK2 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

	20°C stars immediately of a sur-		
	20°C storage immediately after use.		
MSDS:	Download		
Publication Ref	erence		
novel progno Liu CH, Chen Cheng HC, C	f protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma. TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, hen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 013 Feb 8. [Epub ahead of print]		
Applications			
In situ Proximity	Ligation Assay (Cell)		
MAP2K6 TAOK2			
Gene Informatio	on		
Entrez GenelD:	<u>5608</u>		
Gene Name:	MAP2K6		
Gene Alias:	MAPKK6,MEK6,MKK6,PRKMK6,SAPKK3		
Gene Description:	mitogen-activated protein kinase kinase 6		
Omim ID:	<u>601254</u>		
Gene Ontology	: <u>Hyperlink</u>		
Gene Summary	This gene encodes a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis. [provided by RefSeq		
Other Designations:	protein kinase, mitogen-activated, kinase 6 (MAP kinase kinase 6)		
Gene Informatio	on		
Entrez GenelD:	<u>9344</u>		
Gene Name:	TAOK2		
Gene Alias:	KIAA0881,MAP3K17,PSK,PSK1,TAO1,TAO2		
Gene Description:	TAO kinase 2		
Gene Ontology	: <u>Hyperlink</u>		
Other Designations:	prostate derived STE20-like kinase PSK,thousand and one amino acid protein kinase		

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.