



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



RICTOR & AKT1 Protein Protein Interaction Antibody Pair

Catalog # : DI0520

規格 : [1 Set]


List All

Specification

Product Description: This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the RICTOR protein, and the other against the AKT1 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

Application Image


In situ Proximity Ligation Assay (Cell)



Reactivity: Human

Quality Control Protein protein interaction immunofluorescence result.

Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between RICTOR and AKT1. HeLa cells were stained with anti-RICTOR rabbit purified polyclonal antibody 1:1200 and anti-AKT1 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware ([BlobFinder](#)) download from The Centre for Image Analysis at Uppsala University.

Supplied Product: Antibody pair set content:

1. RICTOR rabbit purified polyclonal antibody (20 ug)
2. AKT1 mouse monoclonal antibody (40 ug)

*Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

20°C storage immediately after use.

MSDS:

[!\[\]\(c507f772dba2b921f86777f01218e570_img.jpg\) Download](#)

Publication Reference

1. [An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma.](#)

Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Jiang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

Applications

In situ Proximity Ligation Assay (Cell)

[AKT1 RICTOR](#)

Gene Information

Entrez GeneID: [253260](#)

Gene Name: RICTOR

Gene Alias: DKFZp686B11164, KIAA1999, MGC39830, mAVO3

Gene: rapamycin-insensitive companion of mTOR

Description:

Omim ID: [609022](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: RICTOR and MTOR (FRAP1; MIM 601231) are components of a protein complex that integrates nutrient- and growth factor-derived signals to regulate cell growth (Sarbassov et al., 2004 [PubMed 15268862]).
[supplied by OMIM]

Other: TORC2-specific protein AVO3,pianissimo

Designations:

Gene Information

Entrez GeneID: [207](#)

Gene Name: AKT1

Gene Alias: AKT, MGC99656, PKB, PKB-ALPHA, PRKBA, RAC, RAC-ALPHA

Gene: v-akt murine thymoma viral oncogene homolog 1

Description:

Omim ID: [164730, 181500](#)


Gene Ontology: [Hyperlink](#)

Gene Summary: The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of

the apoptotic machinery. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq]

Other Designations: RAC-alpha serine/threonine-protein kinase,murine thymoma viral (v-akt) oncogene homolog-1,protein kinase B,rac protein kinase alpha

Interactome



[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.