



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## CDKN1A & SKP2 Protein Protein Interaction Antibody Pair

Catalog # : DI0568

規格 : [ 1 Set ]

List All

### Specification


#### Product

#### Description:

This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the CDKN1A protein, and the other against the SKP2 protein for use in *in situ* Proximity Ligation Assay. See Publication Reference below.

### Application Image

*In situ* Proximity Ligation Assay  
(Cell)




#### Reactivity:

Human

#### Quality Control

Protein protein interaction immunofluorescence result.

#### Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between CDKN1A and SKP2. HeLa cells were stained with anti-CDKN1A rabbit purified polyclonal antibody 1:1200 and anti-SKP2 mouse purified polyclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware ([BlobFinder](#)) download from The Centre for Image Analysis at Uppsala University.

#### Supplied

#### Product:

Antibody pair set content:

1. CDKN1A rabbit purified polyclonal antibody (20 ug)

2. SKP2 mouse purified polyclonal antibody (40 ug)

\*Reagents are sufficient for at least 30-50 assays using recommended protocols.

#### Storage

#### Instruction:

Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

**MSDS:****Publication Reference**

1. An analysis of protein-protein interactions in cross-talk pathways reveals CRKL as a novel prognostic marker in hepatocellular carcinoma. Liu CH, Chen TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, Cheng HC, Chen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell Proteomics. 2013 Feb 8. [Epub ahead of print]

**Applications*****In situ* Proximity Ligation Assay (Cell)**

CDKN1A SKP2

**Gene Information****Entrez GeneID:** [1026](#)**Gene Name:** CDKN1A**Gene Alias:** CAP20,CDKN1,CIP1,MDA-6,P21,SDI1,WAF1,p21CIP1**Gene Description:** cyclin-dependent kinase inhibitor 1A (p21, Cip1)**Omim ID:** [116899](#)**Gene Ontology:** [Hyperlink](#)

**Gene Summary:** This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen (PCNA), a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation. Two alternatively spliced variants, which encode an identical protein, have been reported. [provided by RefSeq]

**Other Designations:** CDK-interaction protein 1,DNA synthesis inhibitor,OTTHUMP00000016298,cyclin-dependent kinase inhibitor 1A,melanoma differentiation associated protein 6,wild-type p53-activated fragment 1

**Gene Information****Entrez GeneID:** [6502](#)**Gene Name:** SKP2**Gene Alias:** FBL1,FBXL1,FLB1,MGC1366**Gene Description:** S-phase kinase-associated protein 2 (p45)**Omim ID:** [601436](#)

**Gene Ontology:** [Hyperlink](#)

**Gene Summary:** This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains 10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B (CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and interacts with S-phase kinase-associated protein 1 (SKP1 or p19). In addition, this gene is established as a protooncogene causally involved in the pathogenesis of lymphomas. Alternative splicing of this gene generates 2 transcript variants encoding different isoforms. [provided by RefSeq]

<b>Other Designations:</b>	CDK2/cyclin A-associated protein p45,S-phase kinase-associated protein 2
----------------------------	--

## Interactome

