

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



## **BIRC3 & CASP9 Protein Protein Interaction Antibody Pair**

Catalog # : DI0576

規格:[1Set]

List All


Specification				Application Image
Product Description:	This protein protein interaction antibody pair set comes with two antibodies to detect the protein-protein interaction, one against the BIRC3 protein, and the other against the CASP9 protein for use in <u>in</u> <u>situ</u> Proximity Ligation Assay. See Publication Reference below.			<i>In situ</i> Proximity Ligation Assay (Cell)
	1. Incubate with target primary antibodies	2. Add PLA probes PLUS and MINUS	3. Hybridize connector oligos	
	4. Ligation to form a complete DNA circle	5. Rolling circle amplification	6. Add fluorescent probes to reveal interaction	

**Reactivity:** 

Human

Quality Control Protein protein interaction immunofluorescence result.


Testing:



Representative image of Proximity Ligation Assay of protein-protein interactions between BIRC3 and CASP9. HeLa cells were stained with anti-BIRC3 rabbit purified polyclonal antibody 1:1200 and anti-CASP9 mouse monoclonal antibody 1:50. Each red dot represents the detection of protein-protein interaction complex. The images were analyzed using an optimized freeware (BlobFinder) download from The Centre for Image Analysis at Uppsala University.

Supplied Product:	Antibody pair set content: 1. BIRC3 rabbit purified polyclonal antibody (20 ug) 2. CASP9 mouse monoclonal antibody (40 ug) *Reagents are sufficient for at least 30-50 assays using recommended protocols.
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -

	20°C storage immediately after use.	
MSDS:	m Download	
Publication Ref	erence	
novel prognos Liu CH, Chen Cheng HC, Cl	f protein-protein interactions in cross-talk pathways reveals CRKL as a stic marker in hepatocellular carcinoma. TC, Chau GY, Jan YH, Chen CH, Hsu CN, Lin KT, Juang YL, Lu PJ, nen MH, Chang CF, Ting YS, Kao CY, Hsiao M, Huang CY. Mol Cell 013 Feb 8. [Epub ahead of print]	
Applications		
<i>In situ</i> Proximity	Ligation Assay (Cell)	
BIRC3 CASP9		
Gene Informatio	on	
Entrez GenelD:	330	
Gene Name:	BIRC3	
Gene Alias:	AIP1,API2,CIAP2,HAIP1,HIAP1,MALT2,MIHC,RNF49	
Gene Description:	baculoviral IAP repeat-containing 3	
Omim ID:	<u>601721</u>	
Gene Ontology	<u>Hyperlink</u>	
Gene Summary	The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis by binding to tumor necrosis factor receptor- associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. The amino acid sequence predicts three baculovirus IAP repeat domains and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq	
Other Designations:	TNFR2-TRAF signaling complex protein,apoptosis inhibitor 2,baculoviral IAP repeat-containing protein 3,inhibitor of apoptosis protein 1,mammalian IAP homolog C	
Gene Informatio	on	
Entrez GeneID:	842	
Gene Name:	CASP9	
Gene Alias:	APAF-3,APAF3,CASPASE-9c,ICE-LAP6,MCH6	
Gene Description:	caspase 9, apoptosis-related cysteine peptidase	
Omim ID:	602234	
Gene Ontology	Hyperlink	
Gene Summary	This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic	



服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.