



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

ESR1(phospho S167) & ESR1 Protein Phosphorylation Antibody Pair

Catalog Number: DP0158

Regulatory Status: For research use only (RUO)

Product Description: This protein phosphorylation antibody pair set comes with two antibodies, one against the ESR1 protein, and the other against the specific S167 phosphorylated site of ESR1 for use in [in situ Proximity Ligation Assay](#). [See Publication Reference below](#).

Reactivity: Human

Applications: PLA-Ce

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Supplied Product: Antibody pair set content:

1. Phospho-ESR1 S167 rabbit polyclonal antibody (20 ul)
In PBS (without Mg²⁺ and Ca²⁺), 150 mM NaCl, pH 7.4 (0.02% sodium azide, 50% glycerol)
 2. ESR1 mouse monoclonal antibody (40 ug)
In 1x PBS, pH 7.2
- *Reagents are sufficient for at least 30-50 assays using recommended protocols.

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Entrez GeneID: 2099

Gene Symbol: ESR1

Gene Alias: DKFZp686N23123, ER, ESR, ESRA, Era, NR3A1

Gene Summary: This gene encodes an estrogen receptor, a ligand-activated transcription factor

composed of several domains important for hormone binding, DNA binding, and activation of transcription. The protein localizes to the nucleus where it may form a homodimer or a heterodimer with estrogen receptor 2. Estrogen and its receptors are essential for sexual development and reproductive function, but also play a role in other tissues such as bone. Estrogen receptors are also involved in pathological processes including breast cancer, endometrial cancer, and osteoporosis. Alternative splicing results in several transcript variants, which differ in their 5' UTRs and use different promoters. [provided by RefSeq]