



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TAF15 Split FISH Probe

Catalog Number: FS0101

Regulation Status: For research use only (RUO)

Product Description: Labeled FISH probes for identification of gene split using Fluorescent In Situ Hybridization Technique. ([Technology](#))

Probe 1:

Size:

Fluorophore:

Location: TAF15

Approximately 590kb

Texas Red

17q12

Probe 2:

Size:

Fluorophore:

Location: TAF15

Approximately 610kb

FITC

17q12

Regulatory Status: For research use only (RUO)

Source: Genomic DNA

Origin: Human

Notice: We **strongly recommend** the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: [KA2375](#) or [KA2691](#)) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Supplied Product: DAPI Counterstain (1500 ng/mL)
125 uL for each 100 uL FISH Probe

Storage Instruction: Store at 4°C in the dark.

Entrez GeneID: 8148

Gene Symbol: TAF15

Gene Alias: Npl3, RBP56, TAF2N, TAFII68, hTAFII68

Gene Summary: Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a subunit of TFIID present in a subset of TFIID complexes. Translocations involving chromosome 17 and chromosome 9, where the gene for the nuclear receptor CSMF is located, result in a gene fusion product that is an RNA binding protein associated with a subset of extraskelatal myxoid chondrosarcomas. Two transcripts encoding different isoforms have been identified. [provided by RefSeq]