



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### ERBB2 Split FISH Probe

**Catalog Number:** FS0109

**Regulation Status:** For research use only (RUO)

**Product Description:** Labeled FISH probes for identification of gene split using Fluorescent In Situ Hybridization Technique. ([Technology](#))

**Probe 1:**

**Size:**

**Fluorophore:**

**Location:** ERBB2

Approximately 580kb

Texas Red

17q12

**Probe 2:**

**Size:**

**Fluorophore:**

**Location:** ERBB2

Approximately 790kb

FITC

17q12

**Regulatory Status:** For research use only (RUO)

**Source:** Genomic DNA

**Origin:** Human

**Notice:** We **strongly recommend** the customer to use FFPE FISH PreTreatment Kit 1 (Catalog #: [KA2375](#) or [KA2691](#)) for the pretreatment of Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections.

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Supplied Product:** DAPI Counterstain (1500 ng/mL )  
125 uL for each 100 uL FISH Probe

**Storage Instruction:** Store at 4°C in the dark.

**Entrez GeneID:** 2064

**Gene Symbol:** ERBB2

**Gene Alias:** CD340, HER-2, HER-2/neu, HER2, NEU, NGL, TKR1

**Gene Summary:** This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq]