



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NAT2 (Human) IP-WB Antibody Pair

Catalog # : H00000010-PW1

規格 : [1 Set]


List All

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of NAT2 transfected lysate using rabbit polyclonal anti-NAT2 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with mouse purified polyclonal anti-NAT2.

Supplied Product: Antibody pair set content:
 1. Antibody pair for IP: rabbit polyclonal anti-NAT2 (300 ul)
 2. Antibody pair for WB: mouse purified polyclonal anti-NAT2 (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

MSDS:  [Download](#)

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [10](#)

Gene Name: NAT2

Gene Alias: AAC2,PNAT

Gene Description: N-acetyltransferase 2 (arylamine N-acetyltransferase)

Omim ID: [243400](#)

Gene Ontology: [Hyperlink](#)

Application Image

Immunoprecipitation-Western Blot

Gene Summary: This gene encodes an enzyme that functions to both activate and deactivate arylamine and hydrazine drugs and carcinogens. Polymorphisms in this gene are responsible for the N-acetylation polymorphism in which human populations segregate into rapid, intermediate, and slow acetylator phenotypes. Polymorphisms in this gene are also associated with higher incidences of cancer and drug toxicity. A second arylamine N-acetyltransferase gene (NAT1) is located near this gene (NAT2). [provided by RefSeq]

Other Designations: N-acetyltransferase 2, arylamide acetylase 2, arylamine N-acetyltransferase 2

Gene Pathway

[Caffeine metabolism](#) [Drug metabolism - other enzymes](#) [Metabolic pathways](#)

Related Disease

[Abnormalities, Drug-Induced](#) [Acute Disease](#) [Adenocarcinoma](#) [Adenoma](#)
[Adenomatous Polyposis Coli](#) [Alveolar Bone Loss](#) [Alzheimer Disease](#) [Alzheimer disease](#)
[Aneuploidy](#) [Arthritis, Rheumatoid](#) [Asthma](#) [Asthma](#) [Astrocytoma](#) [Atherosclerosis](#)
[Behcet Syndrome](#) [Birth Weight](#) [Brain Neoplasms](#) [Breast cancer](#) [Breast Neoplasms](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.