



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Datasheet

ARRB2 monoclonal antibody (MP01J), clone 3G1 (PE)

Catalog Number: H00000409-MP01J

Regulation Status: For research use only (RUO)

Product Description: PE conjugated mouse monoclonal antibody raised against a partial recombinant ARRB2.

This product is belong to Cell Culture Grade Antibody (CX Grade).

Clone Name: 3G1

Immunogen: ARRB2 (AAH07427.1, 300 a.a. ~ 409 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence:

NLASSTIVKEGANKEVLGILVSYRVKVKLVSRGGDVS
VELPFVLMHPKPHDHIPLRPQSAAPETDVPVDTNLIE
FDTNYATDDDIVFEDFARLRLKGMKDDDYDDQLC

Host: Mouse

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: PE

Preparation Method: Cell Culture Production

Isotype: IgG2a Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to
avoid repeated freezing and thawing.

Entrez GeneID: 409

Gene Symbol: ARRB2

Gene Alias: ARB2, ARR2, BARR2, DKFZp686L0365

Gene Summary: Members of arrestin/beta-arrestin
protein family are thought to participate in agonist-
mediated desensitization of G-protein-coupled receptors

and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 2, like arrestin beta 1, was shown to inhibit beta-adrenergic receptor function in vitro. It is expressed at high levels in the central nervous system and may play a role in the regulation of synaptic receptors. Besides the brain, a cDNA for arrestin beta 2 was isolated from thyroid gland, and thus it may also be involved in hormone-specific desensitization of TSH receptors. Multiple alternatively spliced transcript variants have been found for this gene, but the full-length nature of some variants has not been defined. [provided by RefSeq]