



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CDA (Human) IP-WB Antibody Pair

Catalog # : H00000978-PW2

規格 : [1 Set]


List All

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of CDA transfected lysate using rabbit polyclonal anti-CDA and Protein A Magnetic Bead (U0007), and immunoblotted with mouse purified polyclonal anti-CDA.

Supplied Product: Antibody pair set content:
 1. Antibody pair for IP: rabbit polyclonal anti-CDA (300 ul)
 2. Antibody pair for WB: mouse purified polyclonal anti-CDA (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [978](#)

Gene Name: CDA

Gene Alias: CDD

Gene Description: cytidine deaminase

Omim ID: [123920](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes an enzyme involved in pyrimidine salvaging. The

encoded protein forms a homotetramer that catalyzes the irreversible hydrolytic deamination of cytidine and deoxycytidine to uridine and deoxyuridine, respectively. It is one of several deaminases responsible for maintaining the cellular pyrimidine pool. Mutations in this gene are associated with decreased sensitivity to the cytosine nucleoside analogue cytosine arabinoside used in the treatment of certain childhood leukemias. [provided by RefSeq]

Other Designations: OTTHUMP0000002883,cytidine aminohydrolase,cytosine nucleoside deaminase,small cytidine deaminase

Gene Pathway

[Drug metabolism - other enzymes](#) [Metabolic pathways](#) [Pyrimidine metabolism](#)

Related Disease

[Adenocarcinoma](#) [Ataxia telangiectasia](#) [Carcinoma, Non-Small-Cell Lung](#) [Carcinoma, Pancreatic Ductal](#) [Colorectal Neoplasms](#) [Disease Progression](#) [Esophageal Neoplasms](#) [Genetic Predisposition to Disease](#) [Hearing Loss](#) [Kidney Failure, Chronic](#) [Leukemia](#) [Leukemia, Myelocytic, Acute](#) [Leukemia, Myeloid, Acute](#) [Lung Neoplasms](#) [Neoplasm Recurrence, Local](#) [Neoplasms](#) [Pancreatic Neoplasms](#) [Recurrence](#) [Stomach Neoplasms](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.