



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CTH monoclonal antibody (MP02J), clone 2E12-1C10 (PE)

Catalog Number: H00001491-MP02J

Regulation Status: For research use only (RUO)

Product Description: PE conjugated mouse monoclonal antibody raised against a full-length recombinant CTH.
This product is belong to Cell Culture Grade Antibody (CX Grade).

Clone Name: 2E12-1C10

Immunogen: CTH (AAH15807, 1 a.a. ~ 405 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence:

MQEKDASSQGFLPHFQHFATQAIHVGQDPEQWTSRA
VVPPISLSTTFKQGAPGQHSGFEYSRSGNPTRNCLEK
AVAALDGAKYCLAFASGLAATVTITHLLKAGDQIICMDD
VYGGTNRYFRQVASEFGLKISFVDCSKIKLLEAAITPET
KLWVIETPTNPTQKVIDIEGCAHIVHKHGDIIILVDNTFM
SPYFQRPLALGADISMYSATKYMNGRSDVVMGLVSVN
CESLHNRLRFLQNSLGAVPSPIDCYLCNRGLKTLHVR
MEKHFKNGMAVAQFLESNPWVEKVIYPGLPSHPQHEL
VKRQCTGCTGMVTFYIKGTLQHAEIFLKNLKLFTLAESL
GGFESLAELPAIMTHASVLKNDRDVLGISDTLIRLSVGL
EDEEDLLEDLDQALKAHPPSGSHS

Host: Mouse

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: PE

Preparation Method: Cell Culture Production

Isotype: IgG1 Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to
avoid repeated freezing and thawing.

Entrez GeneID: 1491

Gene Symbol: CTH

Gene Alias: MGC9471

Gene Summary: This gene encodes a cytoplasmic enzyme in the trans-sulfuration pathway that converts cystathione derived from methionine into cysteine. Glutathione synthesis in the liver is dependent upon the availability of cysteine. Mutations in this gene cause cystathioninuria. Alternative splicing of this gene results in two transcript variants encoding different isoforms. [provided by RefSeq]