

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





CYP3A4 (Human) IP-WB Antibody Pair

Catalog #: H00001576-PW1 規格:[1 Set]

List All

Specification		Application Image
Product Description:	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.	Immunoprecipitation-Wester Blot
Reactivity:	Human	
Quality Control Testing:	Immunoprecipitation-Western Blot (IP-WB) 258 25- 20- 15-	
	Immunoprecipitation of CYP3A4 transfected lysate using rabbit polyclonal anti-CYP3A4 and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with mouse purified polyclonal anti-CYP3A4.	
Supplied Product:	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-CYP3A4 (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-CYP3A4 (50 ug)	
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.	
MSDS:	<u>Download</u>	
Applications		
Immunoprecipit	ation-Western Blot	
Gene Information	on	
Entrez GeneID:	<u>1576</u>	
Gene Name:	CYP3A4	
Gene Alias:	CP33,CP34,CYP3A,CYP3A3,HLP,MGC126680,NF- 25,P450C3,P450PCN1	
Gene Description:	cytochrome P450, family 3, subfamily A, polypeptide 4	
Omim ID:	<u>124010</u>	

Gene Ontology: Hyperlink

Gene Summary: This gene, CYP3A4, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by glucocorticoids and some pharmacological agents. This enzyme is involved in the metabolism of approximately half the drugs which are are used today, including acetaminophen, codeine, cyclosporin A, diazepam and erythromycin. The enzyme also metabolizes some steroids and carcinogens. This gene is part of a cluster of cytochrome P450 genes on chromosome 7q21.1. Previously another CYP3A gene, CYP3A3, was thought to exist; however, it is now thought that this sequence represents a transcript variant of CYP3A4. [provided by RefSeq

Other Designations:

P450-III, steroid inducible, cytochrome P450, subfamily IIIA (niphedipine oxidase), polypeptide 3,cytochrome P450, subfamily IIIA (niphedipine oxidase), polypeptide 4,cytochrome P450, subfamily IIIA, polypeptide 4, glucocorticoid-inducible P450, nifedipine o

Gene Pathway

<u>Drug metabolism - cytochrome P450</u> <u>Drug metabolism - other enzymes</u> gamma-Hexachlorocyclohexane degradation Linoleic acid metabolism Metabolic pathways Metabolism of xenobiotics by cytochrome P450 Retinol metabolism

Related Disease

Acquired Immunodeficiency Syndrome Acute Disease Adenocarcinoma Adenoma, Liver Cell Akathisia, Drug-Induced Anemia, Anemia, Aplastic Asthma Asthma Balkan Nephropathy Body Weight Breast cancer Breast Neoplasms Carcinoma Carcinoma, Ductal, Breast Carcinoma, Hepatocellular Carcinoma, Non-Small-Cell Lung Carcinoma, Renal Cell Carcinoma, Small Cell

... see more

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 2 of 2 2016/5/20