



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DUT (Human) IP-WB Antibody Pair

Catalog # : H00001854-PW2

規格 : [1 Set]


List All

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of DUT transfected lysate using rabbit polyclonal anti-DUT and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with mouse purified polyclonal anti-DUT.

Supplied Product: Antibody pair set content:
 1. Antibody pair for IP: rabbit polyclonal anti-DUT (300 ul)
 2. Antibody pair for WB: mouse purified polyclonal anti-DUT (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [1854](#)

Gene Name: DUT

Gene Alias: FLJ20622,dUTPase

Gene Description: deoxyuridine triphosphatase

Omim ID: [601266](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes an essential enzyme of nucleotide metabolism. The

encoded protein forms a ubiquitous, homotetrameric enzyme that hydrolyzes dUTP to dUMP and pyrophosphate. This reaction serves two cellular purposes: providing a precursor (dUMP) for the synthesis of thymine nucleotides needed for DNA replication, and limiting intracellular pools of dUTP. Elevated levels of dUTP lead to increased incorporation of uracil into DNA, which induces extensive excision repair mediated by uracil glycosylase. This repair process, resulting in the removal and reincorporation of dUTP, is self-defeating and leads to DNA fragmentation and cell death. Alternative splicing of this gene leads to different isoforms that localize to either the mitochondrion or nucleus. A related pseudogene is located on chromosome 19. [provided by RefSeq

Other Designations: dUTP nucleotidohydrolase,dUTP pyrophosphatase,deoxyuridine 5'-triphosphate nucleotidohydrolase

Gene Pathway

[Metabolic pathways](#) [Pyrimidine metabolism](#)

Related Disease

[DNA Damage](#) [Genetic Predisposition to Disease](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.