

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





E2F6 (Human) IP-WB Antibody Pair

Catalog #: H00001876-PW1 規格:[1 Set]

List All

Specification		Application Image
Product Description:	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.	Immunoprecipitation-Western Blot
Reactivity:	Human	
Quality Control Testing:	Immunoprecipitation-Western Blot (IP-WB) 250 -	
	Immunoprecipitation of E2F6 transfected lysate using rabbit polyclonal anti-E2F6 and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with mouse polyclonal anti-E2F6.	
Supplied Product:	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-E2F6 (300 ul) 2. Antibody pair for WB: mouse polyclonal anti-E2F6 (50 ul)	
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.	
MSDS:	<u>Download</u>	
Applications		
Immunoprecipit	ation-Western Blot	
Gene Information	on	
Entrez GeneID:	1876	
Gene Name:	E2F6	
Gene Alias:	E2F-6,MGC111545	
Gene Description:	E2F transcription factor 6	
Omim ID:	602944	
Gene Ontology:		

Page 1 of 2 2016/5/20

Gene Summary: This gene encodes a member of the E2F transcription factor protein family. E2F family members play a crucial role in control of the cell cycle and of the action of tumor suppressor proteins. They are also a target of the transforming proteins of small DNA tumor viruses. Many E2F proteins contain several evolutionarily conserved domains: a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. The encoded protein of this gene is atypical because it lacks the transactivation and tumor suppressor protein association domains. It contains a modular suppression domain and is an inhibitor of E2F-dependent transcription. The protein is part of a multimeric protein complex that contains a histone methyltransferase and the transcription factors Mga and Max. Multiple transcript variants have been reported for this gene, but it has not been clearly demonstrated that they encode valid isoforms. [provided by RefSeq

Other

E2F transcription factor 6, isoform 1

Designations:

Related Disease

Genetic Predisposition to Disease Ovarian Neoplasms

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 2 of 2 2016/5/20