



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

ESR2 (Human) Recombinant Protein (P01)

Catalog Number: H00002100-P01

Regulation Status: For research use only (RUO)

Product Description: Human ESR2 full-length ORF (AAH24181, 1 a.a. - 323 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MDIKNSPSSLNSPSSYNCSQSILPLEHGSIIYIPSSYVDS
HHEYPAAMTFYSPAVMNYISPSNVNLEGGPGRQTTSP
NVLWPTPGHLSPLVHRQLSHLYAEPQKSPRCEARSL
EHTLPVNRETLKRKVSNGRCASPTGPGSKRDAHFCA
VCSDYASGYHYGVWSCEGCKAFFKRSIQGHNDYICPA
TNQCTIDKNRRKSCQACRLRKCYEVGMVKCGSRER
CGYRLVRRQRSADQLHCAGKAKRSGGHAPRVRELL
LDALSPEQLVLTLLAEPPHVLISRPSAPFTEASMMMS
LTKLADKELVHMISWAKKIPGMRGNA

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 61.27

Interspecies Antigen Sequence: Mouse (87)

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 2100

Gene Symbol: ESR2

Gene Alias: ER-BETA, ESR-BETA, ESRB, ESTRB, Erb, NR3A2

Gene Summary: This gene encodes a member of the family of estrogen receptors and superfamily of nuclear receptor transcription factors. The gene product contains an N-terminal DNA binding domain and C-terminal ligand binding domain and is localized to the nucleus, cytoplasm, and mitochondria. Upon binding to 17beta-estradiol or related ligands, the encoded protein forms homo- or hetero-dimers that interact with specific DNA sequences to activate transcription. Some isoforms dominantly inhibit the activity of other estrogen receptor family members. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been fully characterized. [provided by RefSeq]

References:

1. GR and LSD1/KDM1A-Targeted Gene Activation Requires Selective H3K4me2 Demethylation at Enhancers. Clark EA, Wu F, Chen Y, Kang P, Kaiser UB, Fang R, Shi YG. Cell Rep. 2019 Jun 18;27(12):3522-3532.e3. doi: 10.1016/j.celrep.2019.05.062.
2. A Fragment of Adhesion Molecule L1 Binds to Nuclear Receptors to Regulate Synaptic Plasticity and Motor Coordination. Kraus K, Kleene R, Henis M, Braren I, Kataria H, Sharaf A, Loers G, Schachner M, Lutz D. Mol Neurobiol. 2018 Jan 30. [Epub ahead of print]