



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

FH monoclonal antibody (MF09), clone 3E7 (FITC)

Catalog Number: H00002271-MF09

Regulation Status: For research use only (RUO)

Product Description: FITC conjugated mouse monoclonal antibody raised against a full length recombinant FH.

Clone Name: 3E7

Immunogen: FH (AAH03108, 33 a.a. ~ 510 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence:

VPSFWPPNAARMASQNSFRIEYDTFGELKVPNDKYYG
AQTVRSTMNFKIGGVTERMPVIAFGILKRAAAEVN
QDYGLDPKIANAIMKAADEVAEGKLNDFPLVWQTG
SGTQTNMNVNEVISNRAIEMLGELGSKIPVHPNDHV
NKSQSSNDTFPTAMHIAAAIEVHEVLLPGLQKLHDALD
AKSKEFAQIIKIGRTHQTDAVPLTLGQEFSGYVQVVKY
AMTRIKAAAMPRIYELAAGGTAVGTGLNTRIGFAEKVAA
KVAALTGLPFVTAPNKFEALAAHDALVELSGAMNTTAC
SLMKIANDIRFLGSGPRSGLGELILPENEPGSSIMPGKV
NPTQCEAMTMVAAQVMGNHVAVTVGGSNGHFELNVF
KPMMIKNVLHSARLLGDASVSFTENCVVGIQANTERIN
KLMNESLMLVTALNPHIGYDKAAKIAKTAHKNGSTLKE
TAIELGYLTAEQFDEWVKPKDMLGPK

Host: Mouse

Interspecies Antigen Sequence: Mouse (96); Rat (97)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: FITC

Isotype: IgG2b Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to
avoid repeated freezing and thawing.

Entrez GeneID: 2271

Gene Symbol: FH

Gene Alias: HLRCC, LRCC, MCL, MCUL1

Gene Summary: The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Krebs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form is targeted to the mitochondrion, where the removal of the extension generates the same form as in the cytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutations in this gene can cause fumarase deficiency and lead to progressive encephalopathy. [provided by RefSeq]