



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IL8 (Human) Matched Antibody Pair

Catalog # : H00003576-AP41

規格 : [1 Set]


List All

Specification

Product Description: This antibody pair set comes with matched antibody pair to detect and quantify protein level of human IL8.

Reactivity: Human

Quality Control Testing: Standard curve using recombinant protein as an analyte.




Sandwich ELISA detection sensitivity ranging from 1 ng/ml to 100 ng/ml.

Supplied Product: Antibody pair set content:
 1. Capture antibody: mouse monoclonal anti-IL8, IgG2a Kappa (100 ug)
 2. Detection antibody: biotinylated mouse monoclonal anti-IL8, IgG2a Kappa (50 ug)
 *Reagents are sufficient for at least 3-5 x 96 well plates using recommended protocols.

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

ELISA Pair (Recombinant protein)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [3576](#)

Gene Name: IL8

Gene Alias: CXCL8,GCP-1,GCP1,LECT,LUCT,LYNAP,MDNCF,MONAP,NAF,NAP-1,NAP1

Gene Description: interleukin 8

Omim ID: [146930](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the CXC chemokine family. This chemokine is one of the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions

as a chemoattractant, and is also a potent angiogenic factor. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. This gene and other ten members of the CXC chemokine gene family form a chemokine gene cluster in a region mapped to chromosome 4q. [provided by RefSeq

Other Designations: T cell chemotactic factor, beta-thromboglobulin-like protein, chemokine (C-X-C motif) ligand 8, emoctakin, granulocyte chemotactic protein 1, lymphocyte-derived neutrophil-activating factor, monocyte-derived neutrophil chemotactic factor, neutrophil-activating p

Gene Pathway

[Bladder cancer](#) [Chemokine signaling pathway](#) [Cytokine-cytokine receptor interaction](#)
[Epithelial cell signaling in Helicobacter pylori infection](#) [Pathways in cancer](#)
[Toll-like receptor signaling pathway](#)

Related Disease

[Acute Disease](#) [Adenocarcinoma](#) [Adenomatous Polyps](#) [Alcoholism](#) [Alzheimer Disease](#)
[Alzheimer disease](#) [Appetite](#) [Arthritis, Juvenile Rheumatoid](#) [Arthritis, Psoriatic](#)
[Arthritis, Rheumatoid](#) [Asthma](#) [Asthma](#) [Atherosclerosis](#) [Atherosclerosis](#) [Atrophy](#)
[Bacterial Infections](#) [Barrett Esophagus](#) [Behcet Syndrome](#) [Biliary Tract Neoplasms](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.