

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





## KNG1 (Human) IP-WB Antibody Pair

**Catalog #**: H00003827-PW2 規格:[1 Set]

#### List All

Specification		Application Image
Product Description:	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.	Immunoprecipitation-Western Blot
Reactivity:	Human	
Quality Control Testing:	Immunoprecipitation-Western Blot (IP-WB)	
	250 = 150 = 100 = 75 =	
	50	
	37 -	
	25= 20=	
	15=	
	10-	
	Immunoprecipitation of KNG1 transfected lysate using mouse monoclonal anti-KNG1 and Protein A Magnetic Bead ( <u>U0007</u> ), and immunoblotted with rabbit polyclonal anti-KNG1.	
Supplied Product:	Antibody pair set content:  1. Antibody pair for IP: mouse monoclonal anti-KNG1 (300 ug)  2. Antibody pair for WB: rabbit polyclonal anti-KNG1 (50 ul)	
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.	
MSDS:	<u>Download</u>	
Applications		
mmunoprecipit	ation-Western Blot	
Gene Information	on	
Entrez GeneID:	3827	
Gene Name:	KNG1	
Gene Alias:	BDK,KNG	
Gene Description:	kininogen 1	
Omim ID:	228960	
Gene Ontology:		

Page 1 of 2 2016/5/21

Gene Summary: High molecular weight kininogen (HMWK) plays an important role in assembly of the plasma kallikrein (see MIM 147910)-kinin system. The KNG1 gene generates both HMWK and low molecular weight kininogen (LMWK) through alternative splicing. Both HMWK and LMWK contain an identical heavy chain consisting of protein domains 1, 2, and 3. However, HMWK contains a 56-kD light chain that consists of domains 5 and 6H, whereas LMWK contains a unique 4-kD light chain that consists of domain 5L. In both proteins, the heavy and light chains are linked by domain 4, which contains the bradykinin (BK) nonapeptide. BK, which is released by plasma kallikrein, is a potent inflammatory mediator that causes vasodilation and enhanced capillary permeability, induces pain, and stimulates production of nitric oxide and prostacyclin (see MIM 601699) from endothelial cells. During vascular damage, BK stimulates smooth muscle proliferation and intimal hypertrophy. Release of BK from HMWK generates a 2-chain HMWK, termed HMWKa, containing the heavy and light chains joined by a disulfide bond (Merkulov et al., 2008 [PubMed 18000168]).[supplied by OMIM

Other

alpha-2-thiol proteinase inhibitor, bradykinin

**Designations:** 

#### Gene Pathway

Complement and coagulation cascades

#### **Related Disease**

Arrhythmias, Cardiac Blood Coagulation Disorders, Inherited Cardiovascular Diseases Coronary Artery Disease Coronary Disease Death, Sudden, Cardiac <u>Diabetes Mellitus, Type 1 Diabetes Mellitus, Type 2 Diabetic Nephropathies Edema</u> Genetic Predisposition to Disease Hypertension Mental Disorders Obesity Obesity Thrombosis Tobacco Use Disorder

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 2 of 2 2016/5/21