



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

PCNA monoclonal antibody (MA02), clone 1G7 (APC)

Catalog Number: H00005111-MA02

Regulation Status: For research use only (RUO)

Product Description: APC conjugated mouse monoclonal antibody raised against a full length recombinant PCNA.

Clone Name: 1G7

Immunogen: PCNA (AAH00491, 1 a.a. ~ 261 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Sequence:

MFEARLVQGSILKKVLEALKDLNEACWDISSSGVNLQ
SMDSSHVSLVQLTLRSEGFDTYRCDRLAMGVNLTS
MSKILKCAGNEDIITLRAEDNADTLALVFEAPNQEKVSD
YEMKLMDLDVEQLGIPEQEYSCVVKMPSGEFARICRD
LSHIGDAVVISCAKDGVKFSASGELGNGNIKLSQTSNV
DKEEEAVTIEMNEPVQLTFALRYLNFFTKATPLSSTVTL
SMSADVPLVVEYKIADMGHLKYYLAPKIEDEEGS

Host: Mouse

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Conjugation: APC

Isotype: IgG2a kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to
avoid repeated freezing and thawing.

Entrez GeneID: 5111

Gene Symbol: PCNA

Gene Alias: MGC8367

Gene Summary: The protein encoded by this gene is
found in the nucleus and is a cofactor of DNA
polymerase delta. The encoded protein acts as a

homotrimer and helps increase the processivity of
leading strand synthesis during DNA replication. In
response to DNA damage, this protein is ubiquitinated
and is involved in the RAD6-dependent DNA repair
pathway. Two transcript variants encoding the same
protein have been found for this gene. Pseudogenes of
this gene have been described on chromosome 4 and
on the X chromosome. [provided by RefSeq]