



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PLAUR (Human) Matched Antibody Pair

Catalog # : H00005329-AP51

規格 : [1 Set]


[List All](#)

Specification

Product Description: This antibody pair set comes with matched antibody pair to detect and quantify protein level of human PLAUR.

Reactivity: Human

Quality Control Testing: Standard curve using PLAUR 293T overexpression lysate (non-denatured) as an analyte.



Sandwich ELISA detection sensitivity ranging from approximately 729x to 3x dilution of the PLAUR 293T overexpression lysate (non-denatured).

Supplied Product: Antibody pair set content:
 1. Capture antibody: mouse monoclonal anti-PLAUR (100 ug)
 2. Detection antibody: rabbit purified polyclonal anti-PLAUR (50 ug)
 *Reagents are sufficient for at least 3-5 x 96 well plates using recommended protocols.

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

ELISA Pair (Transfected lysate)

 [Protocol Download](#)

Gene Information

Entrez GeneID: [5329](#)

Gene Name: PLAUR

Gene Alias: CD87, UPAR, URKR

Gene Description: plasminogen activator, urokinase receptor

Omim ID: [173391](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-

surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage reactions that have not yet been fully defined. [provided by RefSeq]

Other Designations: 2 monocyte activation antigen Mo3,u-plasminogen activator receptor form

Gene Pathway

[Complement and coagulation cascades](#)

Related Disease

[Asthma](#) [Asthma](#) [Bronchial Hyperreactivity](#) [Carcinoma, Non-Small-Cell Lung](#) [Cardiovascular Diseases](#) [Chorioamnionitis](#) [Diabetes Mellitus, Type 2](#) [Edema](#) [Fetal Membranes, Premature Rupture](#) [Genetic Predisposition to Disease](#) [Kidney Failure, Chronic](#) [Lung Neoplasms](#) [Myocardial Infarction](#) [Obstetric Labor, Premature](#) [Ovarian Neoplasms](#) [Placenta Diseases](#) [Pre-Eclampsia](#) [Premature Birth](#) [Pulmonary Disease, Chronic Obstructive](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.