

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





PSMB8 (Human) IP-WB Antibody Pair

Catalog #: H00005696-PW1 規格:[1 Set]

List All

Specification		Application Image
Product Description:	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.	Immunoprecipitation-Wester Blot
Reactivity:	Human	
Quality Control Testing:		
	250 = 150 = 100 =	
	75 - 50 -	
	37 =	
	25-	
	20-	
	15-	
	Immunoprecipitation of PSMB8 transfected lysate using rabbit polyclonal	
	anti-PSMB8 and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with mouse purified polyclonal anti-PSMB8.	
Supplied	Antibody pair set content:	
Product:	Antibody pair for IP: rabbit polyclonal anti-PSMB8 (300 ul) Antibody pair for WB: mouse purified polyclonal anti-PSMB8 (50 ug)	
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.	
MSDS:	<u>Download</u>	
Applications		
Immunoprecipit	tation-Western Blot	
Gene Information	·	
Entrez GeneID:		
Gene Name:	PSMB8	
Gene Alias:	D6S216,D6S216E,LMP7,MGC1491,PSMB5i,RING10,beta5i	
Gene Description:	proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional peptidase 7)	
Omim ID:	<u>177046</u>	
Gene Ontology:	: <u>Hyperlink</u>	
37	Dage 4 of 0	2

Page 1 of 2 2016/5/21

Gene Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is induced by gamma interferon and this gene product replaces catalytic subunit 3 (proteasome beta 5 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit. Two alternative transcripts encoding two isoforms have been identified; both isoforms are processed to yield the same mature subunit. [provided by RefSeq

Other Designations:

OTTHUMP00000029420,OTTHUMP00000029421,OTTHUMP00000062 981, low molecular weight protein 7, macropain subunit C13, multicatalytic endopeptidase complex subunit C13, protease component

C13, proteasome (prosome, macropain) subunit beta type 8 (large multifunctiona

Gene Pathway

Proteasome

Related Disease

<u>Acute Disease</u> <u>Alveolitis, Extrinsic Allergic Arthritis, Juvenile Rheumatoid Arthritis, Reactive</u> <u>Asthma Bronchiolitis, Viral Brucellosis Cardiovascular Diseases Coronary Disease</u> <u>Dermatitis, Atopic Diabetes Mellitus, Type 1 Diabetes Mellitus, Type 2 Diseases in Twins</u> Edema Genetic Predisposition to Disease Hepatitis B Hepatitis C Hepatitis C, Chronic

... see more

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 2 of 2 2016/5/21