



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

RFC4 (Human) IP-WB Antibody Pair

Catalog # : H00005984-PW2

規格 : [1 Set]


[List All](#)

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of RFC4 transfected lysate using rabbit polyclonal anti-RFC4 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with mouse purified polyclonal anti-RFC4.

Supplied Product: Antibody pair set content:
 1. Antibody pair for IP: rabbit polyclonal anti-RFC4 (300 ul)
 2. Antibody pair for WB: mouse purified polyclonal anti-RFC4 (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [5984](#)

Gene Name: RFC4

Gene Alias: A1,MGC27291,RFC37

Gene Description: replication factor C (activator 1) 4, 37kDa

Omim ID: [102577](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The elongation of primed DNA templates by DNA polymerase delta and

DNA polymerase epsilon requires the accessory proteins proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). RFC, also named activator 1, is a protein complex consisting of five distinct subunits of 140, 40, 38, 37, and 36 kD. This gene encodes the 37 kD subunit. This subunit forms a core complex with the 36 and 40 kDa subunits. The core complex possesses DNA-dependent ATPase activity, which was found to be stimulated by PCNA in an in vitro system. Alternatively spliced transcript variants encoding the same protein have been reported. [provided by RefSeq]

Other A1 37 kDa subunit,RFC 37 kDa subunit,activator 1 37 kDa
Designations: subunit,replication factor C (activator 1) 4 (37kD),replication factor C 4

Gene Pathway

[DNA replication](#) [Mismatch repair](#) [Nucleotide excision repair](#)

Related Disease

[Genetic Predisposition to Disease](#) [Graft vs Host Disease](#) [Multiple Sclerosis](#)
[Urinary Bladder Neoplasms](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.