



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CCL18 (Human) IP-WB Antibody Pair

Catalog # : H00006362-PW1

規格 : [1 Set]


[List All](#)

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of CCL18 transfected lysate using rabbit polyclonal anti-CCL18 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with rabbit polyclonal anti-CCL18.

Supplied Product: Antibody pair set content:
1. Antibody pair for IP: rabbit polyclonal anti-CCL18 (300 ul)
2. Antibody pair for WB: rabbit polyclonal anti-CCL18 (50 ul)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

MSDS:  [Download](#)

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [6362](#)

Gene Name: CCL18

Gene Alias: AMAC-1,AMAC1,CKb7,DC-CK1,DCCK1,MIP-4,PARC,SCYA18

Gene Description: chemokine (C-C motif) ligand 18 (pulmonary and activation-regulated)

Omim ID: [603757](#)

Gene Ontology: [Hyperlink](#)

Application Image

Immunoprecipitation-Western Blot

Gene Summary: This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for naive T cells, CD4+ and CD8+ T cells and nonactivated lymphocytes, but not for monocytes or granulocytes. This chemokine attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes. It may play a role in both humoral and cell-mediated immunity responses. [provided by RefSeq]

Other Designations: CC chemokine ligand 18, alternative macrophage activation-associated CC chemokine 1, chemokine (C-C), dendritic, dendritic cell chemokine 1, macrophage inflammatory protein 4, pulmonary and activation-regulated chemokine, small inducible cytokine A18, small indu

Gene Pathway

[Chemokine signaling pathway](#) [Cytokine-cytokine receptor interaction](#)

Related Disease

[Asthma](#) [Asthma Birth Weight](#) [Bronchiolitis, Viral](#) [Cardiovascular Diseases](#) [Diabetes Mellitus, Type 2](#) [Edema](#) [Genetic Predisposition to Disease](#) [Glioblastoma](#) [Glioma](#) [Infant, Premature, Diseases](#) [Leukemia](#) [Leukemia, Myeloid, Acute](#) [Meningeal Neoplasms](#) [Meningioma](#) [Respiratory Syncytial Virus Infections](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.