



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SDCBP (Human) IP-WB Antibody Pair

Catalog # : H00006386-PW3

規格 : [1 Set]


List All

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of SDCBP transfected lysate using rabbit polyclonal anti-SDCBP and Protein A Magnetic Bead (U0007), and immunoblotted with mouse purified polyclonal anti-SDCBP.

Supplied Product: Antibody pair set content:
1. Antibody pair for IP: rabbit polyclonal anti-SDCBP (300 ul)
2. Antibody pair for WB: mouse purified polyclonal anti-SDCBP (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [6386](#)

Gene Name: SDCBP

Gene Alias: MDA-9,ST1,SYCL,TACIP18

Gene Description: syndecan binding protein (syntenin)

Omim ID: [602217](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene was initially identified as a molecule

linking syndecan-mediated signaling to the cytoskeleton. The syntenin protein contains tandemly repeated PDZ domains that bind the cytoplasmic, C-terminal domains of a variety of transmembrane proteins. This protein may also affect cytoskeletal-membrane organization, cell adhesion, protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic reticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq

Other melanoma differentiation associated protein-9,pro-TGF-alpha
Designations: cytoplasmic domain-interacting protein 18,scaffold protein
Pbp1,syntenin

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.