

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



XBP1 (Human) IP-WB Antibody Pair		
Catalog #: H00007494-PW1		
List All		
Specification		Application Image
Product Description:	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.	Immunoprecipitation-Western Blot
Reactivity:	Human	
Quality Control Testing:	Immunoprecipitation-Western Blot (IP-WB)	
Supplied Product:	Antibody pair set content: 1. Antibody pair for IP: rabbit polyclonal anti-XBP1 (300 ul) 2. Antibody pair for WB: mouse purified polyclonal anti-XBP1 (50 ug)	
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.	
MSDS:	Download	
Applications		
Immunoprecipit	ation-Western Blot	
Gene Informatio	on	
Entrez GenelD:	7494	
Gene Name:	XBP1	
Gene Alias:	TREB5,XBP2	
Gene Description:	X-box binding protein 1	
Omim ID:	<u>125480, 194355</u>	
Gene Ontology	Hyperlink Bogo 1 of 2	2016/

Gene Summary: This gene encodes a transcription factor that regulates MHC class II genes by binding to a promoter element referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRNA of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XBP1(S), which is the functionally active transcription factor. The isoform encoded by the unspliced mRNA, XBP1(U), is constitutively expressed, and thought to function as a negative feedback regulator of XBP1(S), which shuts off transcription of target genes during the recovery phase of ER stress. A pseudogene of XBP1 has been identified and localized to chromosome 5. [provided by RefSeq

Other Designations: OTTHUMP00000028682,X-box-binding protein-1

## Related Disease

Amphetamine-Related Disorders Arousal Atherosclerosis Atherosclerosis Bipolar Disorder Depressive Disorder, Major Genetic Predisposition to Disease Hyperhomocysteinemia IgA Deficiency Ischemia Kidney Failure, Chronic Lung Neoplasms Obesity Obesity Personality Assessment Psychiatric Status Rating Scales Pulmonary Disease, Chronic Obstructive Schizophrenia Schizophrenia

... see more

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖 ©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.